

Identification and Estimation of Semi-parametric Censored Dynamic Panel Data Models*

Yingyao Hu and Ji-Liang Shiu

November 2012

Abstract

This study presents a semiparametric identification and estimation method for censored dynamic panel data models and their average partial effects using only two-period data. The proposed method transforms the semi-parametric specification of censored dynamic panel data models into a valid semi-parametric family of PDFs of observables without modeling the distribution of the initial condition. Then the censored dynamic panel data models can be identified by a standard maximum likelihood estimation (MLE). The identifying assumptions are related to the completeness of the families of known semiparametric PDFs corresponding to censored dynamic panel data models and observed conditional density functions between the dependent and explanatory variables. This study shows that the families of PDFs corresponding to dynamic tobit models and dynamic lognormal hurdle models satisfy the identification assumptions with two types of data generating process (DGP). This study proposes a sieve maximum likelihood estimator (sieve MLE) and investigates the finite sample properties of these sieve-based estimators through Monte Carlo analysis. This study presents the dynamic behavior of annual individual health expenditures estimated as an empirical illustration using the dynamic tobit model and data from the Medical Expenditure Panel Survey (MEPS).

Keywords: dynamic lognormal hurdle model, dynamic tobit model, initial condition, nonlinear dynamic panel data model, unobserved covariate, unobserved heterogeneity, endogeneity

*The authors are solely responsible for any remaining errors. The authors can be reached at jishiu@ccu.edu.tw and yhu@jhu.edu.

1. Introduction

The identification and estimation of dynamic panel data models is one of the main challenges in econometrics. These models are appealing in applied research because they consider the lagged value of the dependent variable as one of the explanatory variables and contain observed and unobserved permanent (heterogeneous) or transitory (serially-correlated) individual differences. In dynamic linear panel data models, researchers have developed and compared several instrumental variables (IV) estimators and generalized method of moments (GMM) estimators in the literature (Anderson and Hsiao (1982), Arellano and Bond (1991), Arellano and Bover (1995), Ahn and Schmidt (1995), Kiviet (1995), Blundell and Bond (1998), Hahn (1999), and Hsiao, Hashem Pesaran, and Kamil Tahmiscioglu (2002)).

When the time dimension, T , is fixed in nonlinear panel data models, the presence of the unobserved effect prevents the construction of a log-likelihood function that can be used to estimate structural parameters consistently. This is the so-called incidental parameters problem discussed by Neyman and Scott (1948). However, the dynamic nature of the models leads to the initial conditions problem because integrating the individual unobserved effect out of the distribution raises the issue of how to specify the distribution of the initial condition given unobserved heterogeneity. Wooldridge (2005) proposed finding the distribution conditional on the initial value and the observed history of strictly exogenous explanatory variables to solve the initial conditions problem. Shiu and Hu (2010) adopted the correlated random effect approach for nonlinear dynamic panel data models without specifying the distribution of the initial condition. They used the identification results of the nonclassical measurement error models of Hu and Schennach (2008) to achieve nonparametric identification of nonlinear dynamic panel data model with three periods of data. Honoré (1993), Hu (2002) and Honoré and Hu (2004) used moment restrictions to identify and estimate the parameters of censored dynamic panel data models. Their results were achieved without making distributions of unobserved heterogeneity and the disturbance, but they failed to identify the average partial effects.

Other quantities of interest in nonlinear panel data applications include the partial effects on the mean response, averaged across the population distribution of the unobserved heterogeneity. Chernozhukov, Fernández-Val, Hahn, and Newey (2009) derived bounds for average

effects in nonseparable panel data models and showed that they can tighten considerably for semiparametric discrete choice models. Graham and Powell (2008) studied the average partial effect over the distribution of unobserved heterogeneity, which represents the causal effect of a small change in an endogenous regressor on a continuously-valued outcome of interest. Hoderlein and White (2009) considered identification of distributional effects and average effects in general nonseparable models, allowing for arbitrary dependence between the persistent unobservables and the regressors of interest even if there are only two time periods. However, their approach explicitly rules out lagged dependent variables. Dynamic models focus on the effects of the lagged dependent variables on the current dependent variable, whereas we want to account or control for the influence of all other variables. The effect of a lagged dependent variables reflects the persistence of the dependent variables over time and the amount of this state dependence can be measured by the average partial effect.

This study focuses the identification and estimation of semi-parametric censored dynamic panel data models and their average partial effects with two periods of data. Compared to the identification results in Shiu and Hu (2010), the proposed approach requires a somewhat stronger but places less demand on the time dimension of data. Under a semi-parametric specification, the models are connected to the semi-parametric distribution of Y_{it} conditional on $(X_{it}, Y_{it-1}, U_{it})$, i.e., $f_{Y_{it}|X_{it}, Y_{it-1}, U_{it}; \theta}$ which is called the semi-parametric censored density function. Then, the identification of θ in $f_{Y_{it}|X_{it}, Y_{it-1}, U_{it}; \theta}$ may lead to that of the proposed semi-parametric censored dynamic panel data models. As mentioned earlier, the maximum likelihood estimator (MLE) for the semi-parametric censored density function $f_{Y_{it}|X_{it}, Y_{it-1}, U_{it}; \theta}$ is inconsistent if the time-series dimension is finite and the unobserved covariate U_{it} is correlated with explanatory variables. This study presents an identification strategy similar to the approach in Hu and Shiu (2011a). The identification method transforms the semi-parametric censored density function $f_{Y_{it}|X_{it}, Y_{it-1}, U_{it}; \theta}$ into a valid semi-parametric family of PDFs of observable variables. This identification technique involves three steps of nontrivial transformation associate with the completeness of known PDFs. The first step is to apply the inverse of an integral operator using $f_{Y_{it}|X_{it}, Y_{it-1}, U_{it}; \theta}$ as a kernel. The second step is to integrate out the unobserved covariate. The last step is to normalize the integrated semi-parametric density function created in the second step. The true value of structural parameters can be uniquely determined by maximizing the likelihood function of the transformed semi-parametric fam-

ily of the PDFs of observables. This process also identifies the average partial effect of the censored dynamic panel data models. The nontrivial transformation steps rely on the completeness of the families of known PDFs $f_{Y_{it}|X_{it},Y_{it-1},U_{it};\theta}$ corresponding to censored dynamic panel data models and observed conditional density functions between the dependent and explanatory variables $f_{Y_{it}|X_{it},Y_{it-1},X_{it-1}}$. The completeness assumptions in this study are not restrictive and applicable to certain types of censored dynamic panel data models with common DGPs. Examples include dynamic tobit models and dynamic lognormal hurdle models for the observed conditional density functions $f_{Y_{it}|X_{it},Y_{it-1},X_{it-1}}$ following normal and distributions or the distributions of exponential families.¹

These identification results suggest a semi-parametric sieve maximum likelihood estimator (sieve MLE) for the proposed model. The consistency of the sieve MLE estimator and the asymptotic normality of its parametric components can be directly obtained from the standard treatment in the sieve MLE literature. This study shows how to implement sieve MLE estimators for dynamic tobit models and dynamic lognormal hurdle models. Combining the estimated parametric components with the nuisance parameter for the initial joint distribution makes it possible to derive a consistent estimator for the average partial effect. An apparent advantage of the proposed sieve MLE procedure is that we can estimate these nonlinear dynamic panel data models using two periods of data without specifying initial conditions. This is beneficial because semi-nonparametric estimators usually require a large sample. Another benefit is that the proposed method allows for time dummies, flexible functional forms of state dependence Y_{it-1} such as quadratics or interaction terms, and parametric heteroskedasticity.

The rest of the article is organized as follows. Section 2 presents the identification of censored dynamic panel data models through several nontrivial transformations. Section 3 shows the identification assumptions hold for dynamic tobit models and dynamic lognormal hurdle models for two types of DGPs. Section 4 presents the proposed sieve MLE and inference. Section 5 presents the results of Monte Carlo experiments for dynamic tobit models. Section 6 shows the application of the sieve MLE to a dynamic tobit model describing the dynamic behaviors of annual individual health expenditures using data from the Medical Expenditure Panel Survey (MEPS). Finally, Section 7 provides concluding remarks. Appendices include proofs of each transformation step and a discrete case.

¹More discussions of the completeness condition can be found in D'Haultfoeuille (2011), Andrews (2011), and Hu and Shiu (2011b).

2. Identification of Censored Dynamic Panel Data Models

Suppose $g_1(\cdot, \cdot; \theta_1)$, $g_2(\cdot, \cdot; \theta_2)$ are known parametric functions and g_1 is strictly increasing in its second argument. Consider the following censored dynamic panel data model:

$$Y_{it} = g_1 \left(g_2 \left(X'_{it}, Y_{it-1}; \theta_2 \right), V_i + \varepsilon_{it}; \theta_1 \right), \quad \forall i = 1, \dots, N; t = 1, \dots, T-1, \quad (1)$$

where Y_{it} is the dependent variable, X_{it} is a vector of observed explanatory variables, ε_{it} is a transitory error term, V_i is an unobservable individual-specific effect, and $\theta \equiv (\theta_1, \theta_2)$ is the parameter to be estimated. The functions g_1 and g_2 may be specified by users, such as $g_1(\chi, \nu; \theta_1) = \max(0, \chi + \nu)$ and $g_2(X'_{it}, Y_{it-1}; \theta_2) = X'_{it}\beta + \gamma Y_{it-1}$, etc. The specifications of g_2 can contain time trends, allowing nonlinear relationships such as quadratics or interactions terms. One of the difficulties of distinguishing between structures of Model (1) from observed samples is that the explanatory variables, (X'_{it}, Y_{it-1}, V_i) , and the transitory error term, ε_{it} , are not independently distributed. Assume that a function of variables in the past can purge the statistical dependence that may exist between the explanatory variables and the transitory error term ε_{it} in Model (1).

Assumption 2.1. (*Exogenous Shocks*) Set η_{it} as an unobserved serially-correlated component in the past such that $\eta_{it} = \varphi \left(\{X_{i\tau}, Y_{i\tau-1}, \varepsilon_{i\tau}\}_{\tau=0,1,\dots,t-1} \right)$ for some function φ . Assume that a transitory random shock ξ_{it} is independent of $\{X_{i\tau}, Y_{i\tau-1}, V_i, \varepsilon_{i\tau-1}\}$ for any $\tau \leq t$. Then, the transitory error term ε_{it} has the following decomposition

$$\varepsilon_{it} = \eta_{it} + \xi_{it}. \quad (2)$$

Plugging Eq. (2) into Model (1) leads to

$$\begin{aligned} Y_{it} &= g_1 \left(g_2 \left(X'_{it}, Y_{it-1}; \theta_2 \right), V_i + \eta_{it} + \xi_{it}; \theta_1 \right) \\ &\equiv g_1 \left(g_2 \left(X'_{it}, Y_{it-1}; \theta_2 \right), U_{it} + \xi_{it}; \theta_1 \right), \end{aligned} \quad (3)$$

where $U_{it} = V_i + \eta_{it}$ is an unobserved covariate. To describe every structure of Model (3) by a parameter, assume that the distribution of ξ_{it} has a semi-parametric representation. This

effectively reduces the identification problem to identify a set of parameters.² This framework leads to the following definitions.³

Definition 2.1. Let Θ_α be a parameter space and let $F(\xi; \alpha)$ be a proper distribution function. If $dF(\xi; \alpha_0)$ is the true distribution, then $dF(\xi; \alpha)$ is correctly specified at α_0 . The parameter point α_0 is globally identifiable if there exists no other $\alpha \in \Theta_\alpha$ such that with probability 1, $dF(\xi; \alpha) = dF(\xi; \alpha_0)$, where the measure is taken with respect to α_0 .

Definition 2.2. The parameter point α_0 is locally identifiable if there exists an open neighborhood of α_0 containing no other α such that with probability 1, $dF(\xi; \alpha) = dF(\xi; \alpha_0)$, where the measure is taken with respect to α_0 .

If α_0 is globally identifiable then it is locally identifiable.

Assumption 2.2. (Semi-parametric Distribution) The semi-parametric distribution of the transitory random shock $dF(\xi_{it}; \alpha)$ is known and is correctly specified at an unknown α_0 . The parameter point α_0 is locally identifiable.

Because g_1 is strictly increasing in its second argument, the exogeneity of ξ_{it} makes it possible to obtain

$$F_{\xi_{it}; \alpha}(\xi) = F_{Y_{it}|X_{it}, Y_{it-1}, U_{it}}(g_1(g_2(X'_{it}, Y_{it-1}; \theta_2), U_{it} + \xi; \theta_1) | X_{it}, Y_{it-1}, U_{it}), \quad (4)$$

or

$$g_1(g_2(X'_{it}, Y_{it-1}; \theta_2), U_{it} + \xi; \theta_1) = F_{Y_{it}|X_{it}, Y_{it-1}, U_{it}; \theta}^{-1}(F_{\xi_{it}; \alpha}(\xi) | X_{it}, Y_{it-1}, U_{it}; \theta), \quad (5)$$

if the inverse of $F_{Y_{it}|X_{it}, Y_{it-1}, U_{it}}$ exists and $\theta \equiv (\theta_1, \theta_2, \alpha)'$. Thus, according to Assumptions 2.1 and 2.2, there is a unique conditional distribution associated with each structure in the censored dynamic panel data Model (3) and the identification of the censored dynamic panel data models (1) is implied by that of the distribution of Y_{it} conditional on $(X_{it}, Y_{it-1}, U_{it})$ (i.e., $f_{Y_{it}|X_{it}, Y_{it-1}, U_{it}; \theta}$). Given this semi-parametric representation, the identification problem is to find conditions such that a true underlying parameter θ_0 can be distinguished on the basis of

²The parameters considered are potentially infinite-dimensional ones. We assume a parameter is constituted of two components, a finite-dimensional parameter vector, and a potentially infinite-dimensional nuisance parameter.

³The definitions can be found in Bowden (1973).

sample observations. The conditional PDF $f_{Y_{it}|X_{it},Y_{it-1},U_{it};\theta}$ corresponding to $F_{Y_{it}|X_{it},Y_{it-1},U_{it};\theta}$ is called the semi-parametric censored density function in this paper. We introduce two examples to highlight this important connection. Suppose F and f denote the CDF and the PDF of an independent random shock, respectively.

Example 1 (Dynamic Censored Model with the Lagged Dependent Variable): Assume $g_1(\chi, \nu; \theta_1) = \max(0, \chi + \nu)$.

$$Y_{it} = \max\{0, g_2(X'_{it}, Y_{it-1}; \theta_2) + U_{it} + \xi_{it}\} \quad \text{with} \quad \forall i = 1, \dots, N; t = 1, \dots, T-1. \quad (6)$$

The semi-parametric censored density function is

$$\begin{aligned} f_{Y_{it}|X_{it},Y_{it-1},U_{it};\theta} &= F_{\xi_{it};\alpha}(-g_2(X'_{it}, Y_{it-1}; \theta_2) - U_{it})^{\mathbf{1}(Y_{it}=0)} \\ &\times f_{\xi_{it};\alpha}(Y_{it} - g_2(X'_{it}, Y_{it-1}; \theta_2) - U_{it})^{\mathbf{1}(Y_{it}>0)}. \end{aligned} \quad (7)$$

Example 2 (Dynamic Log Hurdle Model with the Lagged Dependent Variable): Define a binary indicator variable $d_{it} = 1(g_3(X'_{it}, Y_{it-1}; \theta_1) + \varsigma_{it} \geq 0)$ where $1(\cdot)$ is the 0-1 indicator function. Suppose that $Y_{it} > 0$ is observed for $d_{it} = 1$ and $Y_{it} = 0$ for $d_{it} = 0$. When $Y_{it} > 0$,

$$Y_{it} = g_2(X'_{it}, Y_{it-1}; \theta_2) + U_{it} + \xi_{it} \quad \text{with} \quad \forall i = 1, \dots, N; t = 1, \dots, T-1. \quad (8)$$

The conditional distribution of interest is

$$\begin{aligned} f_{Y_{it}|X_{it},Y_{it-1},U_{it};\theta} &= F_{\varsigma_{it}}(-g_3(X'_{it}, Y_{it-1}; \theta_1) - U_{it})^{\mathbf{1}(Y_{it}=0)} \left\{ (1 - F_{\varsigma_{it}}(-g_3(X'_{it}, Y_{it-1}; \theta_1) - U_{it})) \right. \\ &\quad \times f_{\xi_{it};\alpha}(\log(Y_{it}) - g_2(X'_{it}, Y_{it-1}; \theta_2) - U_{it}) \frac{1}{Y_{it}} \left. \right\}^{\mathbf{1}(Y_{it}>0)}. \end{aligned} \quad (9)$$

Many nonlinear dynamic panel data models, such as dynamic discrete choice models, can be converted into their corresponding semi-parametric density functions. However, a discrete dependent variable may incur a very strong restriction that forbids a continuously distributed unobserved covariate. These examples satisfy the identification assumptions when the unobserved covariate is assumed to be continuously distributed.

A number of economic optimization studies have presented empirical applications of these

censored dynamic panel data models, where the dependent variables Y_{it} represents the amount of insurance coverage chosen by an individual, annual women's labor supply, a firm's expenditures on R&D, or annual individual health expenditures. In both examples, the models contain the lagged censored dependent variables in the right-hand side(RHS), ruling out top-coded censored models that contain a lagged value of a latent variable. Because piles of the dependent variable at zero can be regarded as optimal solutions of utility maximizing behavior, these models are often called corner solution models with lagged censored dependent variables.

2.1. General Identification

Consider the semi-parametric censored density function:

$$f_{Y_{it}|X_{it},Y_{it-1},U_{it};\theta}(y_{it}|x_{it},y_{it-1},u_{it}), \quad (10)$$

where Y_{it} is the dependent variable for an individual i , and the explanatory variables include a lagged dependent variable, a set of possibly time-varying explanatory variables X_{it} , and the unobserved covariate U_{it} . As discussed earlier, the identification of this semi-parametric censored density function leads to the identification of certain types of censored dynamic panel data models. Assume that $\theta_0 \in \Theta$ is local identifiable. In other words, θ_0 is a unique value of θ in an open neighborhood of θ_0 , which specifies the exact structure of the model. Consider a panel data containing two periods, $\{Y_{it}, X_{it}, Y_{it-1}, X_{it-1}\}_i$ for $i = 1, 2, \dots, N$. Assume that for each i , $(Y_{it}, X_{it}, Y_{it-1}, X_{it-1})$ is an independent random draw from a bounded distribution $f_{Y_{it}, X_{it}, Y_{it-1}, X_{it-1}}$. The law of total probability leads to the following,

$$f_{Y_{it}, X_{it}, Y_{it-1}, X_{it-1}} = \int f_{Y_{it}|X_{it}, Y_{it-1}, U_{it}; \theta_0} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0} du_{it} \quad (11)$$

where $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0} = f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}}$ is the joint density function of variables $(x_{it}, y_{it-1}, x_{it-1}, u_{it})$. Let \mathcal{Y}_{it} , \mathcal{X}_{it} , and \mathcal{U}_{it} be the support of random variables Y_{it} , X_{it} , and U_{it} , respectively. Set $\mathcal{L}^2(\mathcal{Y}) = \{h(\cdot) : \int_{\mathcal{Y}} |h(y)|^2 dy < \infty\}$ and $\mathcal{L}^2(\mathcal{U}, \omega) = \{h(\cdot) : \int_{\mathcal{U}} |h(u)|^2 \omega(u) du < \infty, \text{ and } \int_{\mathcal{U}} \omega(u) du < \infty\}$. Note the weighted \mathcal{L}^2 -space, $\mathcal{L}^2(\mathcal{U}, \omega)$, contains a constant function (i.e., $c(u) = c \forall u \in \mathcal{U}$). The key idea of this identification method is to extend Eq. (11) to the semi-parametric censored density function (10) over a proper subset of \mathcal{Y}_{it} . The first

step is to construct $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta}$ and avoid an unwanted restriction.⁴ Set Y_{it}^+ as the argument of y_{it} in $\tilde{\mathcal{Y}}_{it} \subsetneq \mathcal{Y}_{it}$

$$\underbrace{f_{Y_{it}^+, X_{it}, Y_{it-1}, X_{it-1}}}_{\text{Observed from Data}} = \int \underbrace{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta}}_{\text{Semi-parametric Specification}} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta} du_{it}. \quad (12)$$

Because the observable density function $f_{Y_{it}^+, X_{it}, Y_{it-1}, X_{it-1}}$ in the LHS and the semi-parametric censored density function $f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta}$ are known, it is possible to construct a semi-parametric joint density function $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta}$ using Eq. (12). Given (x_{it}, y_{it-1}) and a parameter θ , define an integral operator as follows:

$$\begin{aligned} L_{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta}} : \mathcal{L}^2(\mathcal{U}_{it}, \omega) &\rightarrow \mathcal{L}^2(\tilde{\mathcal{Y}}_{it}) \text{ with} \\ (L_{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta}} h)(y_{it}^+) &= \int \frac{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta}(y_{it}^+ | x_{it}, y_{it-1}, u_{it})}{\omega(u_{it})} h(u_{it}) \omega(u_{it}) du_{it}. \end{aligned} \quad (13)$$

If the integral operator $L_{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta}}$ is invertible for each θ , then Eq. (12) suggests that the semi-parametric joint density function can be obtained by

$$f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta} \equiv L_{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta}}^{-1} (f_{Y_{it}^+, X_{it}, Y_{it-1}, X_{it-1}}). \quad (14)$$

Plugging the true parameter θ_0 into this equation results in

$$f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0} \equiv L_{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0}}^{-1} (f_{Y_{it}^+, X_{it}, Y_{it-1}, X_{it-1}})$$

by Eq. (11). The semi-parametric joint density function still achieves the true joint density function at the population parameter θ_0 or it is correctly specified at α_0 . The concept of completeness provides a sufficient condition for the invertibility of the integral operator using the semi-parametric censored density function as a kernel. The following definition presents this completeness.

Definition 2.3. A density function $f(y|u)$ satisfies a completeness condition if for $h(u) \in$

⁴Extend Eq. (11) to θ by $f_{Y_{it}, X_{it}, Y_{it-1}, X_{it-1}} = \int f_{Y_{it} | X_{it}, Y_{it-1}, U_{it}; \theta} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta} du_{it}$ over the whole \mathcal{Y}_{it} . Integrating out y_{it} over \mathcal{Y}_{it} results in $f_{X_{it}, Y_{it-1}, X_{it-1}} = \int f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta} du_{it}$. This suggests that $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta}$ loses the variation over θ after integrating out the observed covariate U_{it} .

$\mathcal{L}^2(\mathcal{U}, \omega)$ such that

$$\int h(u) f(y|u) \omega(u) du = 0 \quad \text{for all } y \quad (15)$$

then $h(u) = 0$ almost everywhere. In other words, there is no nonzero function in $\mathcal{L}^2(\mathcal{U}, \omega)$ with zero integration for each function in the family of the density functions $\{f(y|u) : y \in \mathcal{Y}\}$. By switching the roles of y and u and dropping ω , it is possible to define $\{f(y|u) : u \in \mathcal{U}\}$ as complete in $\mathcal{L}^2(\mathcal{Y})$, and this definition can be generalized to function forms such as $f(y, u)$.

Assumption 2.3. (Dependence between Y_{it} and U_{it}) For each $\theta \in \Theta$ and fixed (x_{it}, y_{it-1}) , the family of the semi-parametric censored density functions $\{\frac{1}{\omega(u_{it})} f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta} : y_{it}^+ \in \tilde{\mathcal{Y}}_{it}\}$ is complete over $\mathcal{L}^2(\mathcal{U}_{it}, \omega)$.

Assumption 2.3 implies that a cardinality restriction in that the cardinality of \mathcal{U}_{it} is less than the cardinality of $\tilde{\mathcal{Y}}_{it}$. Thus, if \mathcal{U}_{it} is a finite discrete set, then the proposed method may apply to a dynamic discrete choice model in which the dependent variable Y_{it} takes more discrete values. However, because of inaccessibility of units of measurement of the unobserved covariate U_{it} , to some extent it is restrictive to assume \mathcal{U}_{it} is discrete.⁵ Therefore, allowing the unobserved covariate U_{it} to take continuous values is more appealing and this study focuses on censored dynamic panel data models.

Suppose that $(L_{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta} h_1})(y_{it}^+) = (L_{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta} h_2})(y_{it}^+)$ for all $y_{it}^+ \in \tilde{\mathcal{Y}}_{it}$. Assumption 2.3 guarantees that $h_1 = h_2$ (i.e., $L_{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta}$ is one-to-one). Hence, the operators $L_{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta}$ are invertible for each θ and (x_{it}, y_{it-1}) . This assumption requires dependence between Y_{it} and U_{it} because the independence between Y_{it} and U_{it} violates Assumption 2.3. Although Assumption 2.3 ensures the existence of the semi-parametric joint density function, $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta}$, it may not be identifiable. The variation of the parameter θ in $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta}$ might be lost in the sense that for any open neighborhood of θ_0 , there exists some $\theta_1 \neq \theta_0$ such that $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_1} = f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}$. In other words, $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta}$ is not locally identifiable. In this case, applying the inverse transformation is useless because the parameter of interest θ is not distinguished in the new transformed joint density functions, preventing the identification of the parameter. The assumption prevents this loss.

⁵The term \mathcal{U}_{it} is a composite error term whose support is involved with the supports of V_i and η_{it} . Assuming discreteness suggests that both V_i and η_{it} are discrete, which may be a strong assumption.

Assumption 2.4. Given each (x_{it}, y_{it-1}) , suppose $f_{X_{it}, Y_{it-1}, X_{it-1}} > 0$. Assume the following conditions:

(i)(Dependence between Y_{it} and X_{it-1}) The family of the observable conditional density functions over \mathcal{X}_{it-1} , $\{f_{Y_{it}^+|X_{it}, Y_{it-1}, X_{it-1}} : x_{it-1} \in \mathcal{X}_{it-1}\}$, is complete over $\mathcal{L}^2(\tilde{\mathcal{Y}}_{it})$.

(ii)(Dependence between Y_{it} and U_{it}) The family of the semi-parametric censored density function over \mathcal{U}_{it} , $\{f_{Y_{it}^+|X_{it}, Y_{it-1}, U_{it}; \theta_0} : u_{it} \in \mathcal{U}_{it}\}$, is complete over $\mathcal{L}^2(\tilde{\mathcal{Y}}_{it})$

This assumption warrants several comments. First, the conditional density functions in the statement are both observable. Second, part (i) suggests that X_{it} cannot be constant over time. If X_{it} is constant across time, then $X_{it} = X_{it-1}$ and $f_{Y_{it}^+|X_{it}, Y_{it-1}, X_{it-1}} = f_{Y_{it}^+|X_{it}, Y_{it-1}}$ which clearly violates the completeness in part (i). Finally, similar to Assumption 2.3, part (ii) requires that, the cardinality of $\tilde{\mathcal{Y}}_{it}$ is less than the cardinality of \mathcal{U}_{it} . Combining the cardinality restrictions in Assumption 2.3 and Assumption 2.4(ii) shows that the cardinality of $\tilde{\mathcal{Y}}_{it}$ is equal to the cardinality of \mathcal{U}_{it} . This restriction is compatible with both the dependent variable Y_{it} and the unobserved covariate U_{it} taking continuous values.

Lemma 2.1. (Applying the inverse) Under Assumptions 2.3-2.4, the semi-parametric joint density $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta}$ is correctly specified at θ_0 and the parameter θ_0 is locally identifiable (i.e., there is an open neighborhood of θ_0 containing no other θ such that $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta} = f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}$).

Proof: See the appendix.

Because $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta}$ contains the unobserved component U_{it} , we need to integrate it out to acquire an observable density function. Set

$$\tilde{f}_{X_{it}, Y_{it-1}, X_{it-1}; \theta}(x_{it}, y_{it-1}, x_{it-1}) \equiv \int f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta}(x_{it}, y_{it-1}, x_{it-1}, u_{it}) du_{it}. \quad (16)$$

To identify θ from the integrated semi-parametric density function $\tilde{f}_{X_{it}, Y_{it-1}, X_{it-1}; \theta}$, it is necessary to examine whether $\tilde{f}_{X_{it}, Y_{it-1}, X_{it-1}; \theta}$ can be correctly specified at θ_0 and the parameter θ_0 is locally identifiable after applying the integration. This integration step might impose too many restrictions on the parameters and degenerate the variation of the function over its parameter space. Thus, it is necessary to rule out these degenerated cases. The following condition maintains the nontrivial semi-parametric representation of $\tilde{f}_{X_{it}, Y_{it-1}, X_{it-1}; \theta}$.

Assumption 2.5. (*Variation of parameters around θ_0*) The family of the derivative of the semi-parametric censored density functions with respect to θ , $\{\frac{\partial}{\partial\theta}f_{Y_{it}^+|X_{it},Y_{it-1},U_{it};\theta_0} : u_{it} \in \mathcal{U}_{it}\}$, is complete over $\mathcal{L}^2(\tilde{\mathcal{Y}}_{it})$.

We summarize the results of the nontrivial transformation of the semi-parametric censored density function after the integration.

Lemma 2.2. (*Integrating out*) Under Assumptions 2.3-2.5, the integrated semi-parametric joint density $\tilde{f}_{X_{it},Y_{it-1},X_{it-1};\theta}$ is correctly specified at θ_0 and the parameter θ_0 is locally identifiable.

Proof: See the appendix.

At this point in the process, the unobserved component of the semi-parametric censored density function (10) has been transformed out and the parameter θ of the function becomes the parameter of the observable semi-parametric function $\tilde{f}_{X_{it},Y_{it-1},X_{it-1};\theta}$. However, if $\tilde{f}_{X_{it},Y_{it-1},X_{it-1};\theta}$ does not integrate to unity (with respect to the measure $dx_{it}dy_{it-1}dx_{it-1}$), it is not a candidate of the semi-parametric family of PDFs for $f_{X_{it},Y_{it-1},X_{it-1}}$ and the standard MLE cannot be applied to $\tilde{f}_{X_{it},Y_{it-1},X_{it-1};\theta}$. To obtain a valid semi-parametric family of PDFs, perform the following normalization step

$$f_{X_{it},Y_{it-1},X_{it-1};\theta} \equiv \frac{\tilde{f}_{X_{it},Y_{it-1},X_{it-1};\theta}}{\int \int \int \tilde{f}_{X_{it},Y_{it-1},X_{it-1};\theta} dx_{it} dy_{it-1} dx_{it-1}}. \quad (17)$$

Similar to the previous discussion, it is necessary to show that the PDF of observables $f_{X_{it},Y_{it-1},X_{it-1};\theta}$ is correctly specified at θ_0 and the parameter θ_0 is locally identifiable after this normalization. The following assumption and lemma demonstrate the existence of a nontrivial θ_0 after normalization.

Assumption 2.6. (*Dependence between Y_{it} and X_{it-1}*) Assume that the family of the observable conditional density functions $\{\frac{\partial}{\partial x_{it-1}}f_{Y_{it}^+|X_{it},Y_{it-1},X_{it-1}} : x_{it-1} \in \mathcal{X}_{it-1}\}$ is complete over $\mathcal{L}^2(\tilde{\mathcal{Y}}_{it})$ for each x_{it}, y_{it-1} .

If \mathcal{X}_{it-1} is discrete, the derivative can be replaced with the difference.⁶ Notice that both Assumption 2.4(i) and Assumption 2.6 are related to the observable conditional distribution

⁶The appendix presents the discussion of a discrete case.

$f_{Y_{it}^+|X_{it},Y_{it-1},X_{it-1}}$ and Assumption 2.6 implies Assumption 2.4(i).⁷ Hence, the two assumptions are compatible and it is only necessary to verify the completeness in Assumption 2.6. The assumption rules out the cases that $f_{Y_{it}^+|X_{it},Y_{it-1},X_{it-1}} = f_1(Y_{it}^+, X_{it}, Y_{it-1})f_2(X_{it-1})$ or $f_{Y_{it}^+|X_{it},Y_{it-1},X_{it-1}} = f_1(Y_{it}^+)f_2(X_{it}, Y_{it-1}, X_{it-1})$.

Lemma 2.3. (*Normalization*) Under Assumptions 2.3, 2.4(ii) and 2.5-2.6, the PDF of observables after normalization, $f_{X_{it},Y_{it-1},X_{it-1};\theta}$, is correctly specified at θ_0 and the parameter θ_0 is locally identifiable.

Proof: See the appendix.

The nontrivial transformation includes applying the inverse of an integral operator using $f_{Y_{it}|X_{it},Y_{it-1},U_{it};\theta}$ as a kernel, integrating out the unobserved covariate, and normalization. After these three steps of transformation associated with the completeness of PDFs, the semi-parametric PDFs of observables $\{f_{X_{it},Y_{it-1},X_{it-1};\theta} : \theta \in \Theta\}$ is correctly specified at θ_0 and the parameter θ_0 is locally identifiable under Assumptions 2.3-2.6. To distinguish the parameters of interest θ_0 from the parameter space Θ on the basis of sample information, use the Kullback-Leibler information criterion

$$K(\theta) = E \left[\log \left(\frac{f_{X_{it},Y_{it-1},X_{it-1};\theta}(x_{it}, y_{it-1}, x_{it-1})}{f_{X_{it},Y_{it-1},X_{it-1}}(x_{it}, y_{it-1}, x_{it-1})} \right) \right] \quad (18)$$

where expectation is taken with respect to $f_{X_{it},Y_{it-1},X_{it-1}}$. Applying the standard framework of the identifiability criterion of maximum likelihood estimation (MLE) produces the following results.

Theorem 2.1. Suppose that $K(\theta) = 0$ has a unique solution at $\theta = \theta_0$ in Θ . Under Assumptions 2.3, 2.4(ii) and 2.5-2.6, the semi-parametric censored density function $f_{Y_{it}|X_{it},Y_{it-1},U_{it};\theta}$ and the joint density function $f_{X_{it},Y_{it-1},X_{it-1},U_{it}}$ can then be identified given the distribution of the two-period observable variables $(y_{it}, x_{it}, y_{it-1}, x_{it-1})$ for $i = 1, 2, \dots, N$.

In addition to the Kullback-Leibler information of classical statistics, the identification result is based on the completeness of the families of known PDFs $f_{Y_{it}|X_{it},Y_{it-1},U_{it};\theta}$ corresponding

⁷Suppose that $h \in \mathcal{L}^2(\tilde{\mathcal{Y}}_{it})$ and $\int h(y_{it}^+) f_{Y_{it}^+|X_{it},Y_{it-1},X_{it-1}} dy_{it}^+ = 0$ for any X_{it-1} . The integration does not involve X_{it-1} and we can take derivative with respect to X_{it-1} . This leads to $\int h(y_{it}^+) \frac{\partial}{\partial x_{it-1}} f_{Y_{it}^+|X_{it},Y_{it-1},X_{it-1}} dy_{it}^+ = 0$ for any X_{it-1} . If $\{\frac{\partial}{\partial x_{it-1}} f_{Y_{it}^+|X_{it},Y_{it-1},X_{it-1}} : x_{it-1} \in \mathcal{X}_{it-1}\}$ satisfies Assumption 2.6, then $h = 0$, which implies that Assumption 2.4(i) holds.

to censored dynamic panel data models and observed conditional density functions between the dependent and explanatory variables $f_{Y_{it}|X_{it},Y_{it-1},X_{it-1}}$. For some semi-parametric specifications, the conditions of completeness are easy to verify or draw inferences from samples. The following sections provide detailed discussions.⁸

Theorem 2.1 provides the identification of the parameter θ . However, because U_{it} does not have meaningful units of measurement, it is not apparent what values of U_{it} we should use. In nonlinear models, estimating the average partial effects of explanatory variables is more attractive than estimating parameters. Thus, this study introduces the average structure function (ASF) by averaging a scalar function of y_{it} , $\omega(y_{it})$, across the distribution of U_{it} in the population. Let (X_{it}, Y_{it-1}) be a given value of the explanatory variables, whose average structure function is

$$\begin{aligned}\mu(X_{it}, Y_{it-1}) &\equiv E_{U_{it}} [E_{Y_{it}} [\omega(y_{it}) | X_{it}, Y_{it-1}, U_{it}]] \\ &= \int_{U_{it}} \left(\int_{Y_{it}} \omega(y_{it}) f_{Y_{it}|X_{it},Y_{it-1},U_{it}} dy_{it} \right) f_{U_{it}} du_{it}.\end{aligned}\quad (19)$$

The marginal distribution of the unobserved covariate U_{it} is also identified by the integration of the joint density function:

$$f_{U_{it}} = \int_{\mathcal{X}_{it}} \int_{\mathcal{Y}_{it-1}} \int_{\mathcal{X}_{it-1}} f_{X_{it},Y_{it-1},X_{it-1},U_{it}} dx_{it} dy_{it-1} dx_{it-1}.\quad (20)$$

Combining the identification results of $f_{Y_{it}|X_{it},Y_{it-1},U_{it}}$ and $f_{U_{it}}$ provides the identification of the average structure function $\mu(X_{it}, Y_{it-1})$. This indicates that the average partial effect is also identified because the average partial effect can be defined by taking derivatives or differences of ASF in Eq. (19) with respect to elements of (X_{it}, Y_{it-1}) . This yields the identification of the average partial effect.

Corollary 2.1. *Under Assumptions 2.3, 2.4(ii) and 2.5-2.6, the average partial effect defined as derivatives or differences of Eq. (19) is identified by a two-period panel data, $\{Y_{it}, X_{it}, Y_{it-1}, X_{it-1}\}$ for $i = 1, 2, \dots, N$.*

The identification condition in Theorem 2.1 that the log-likelihood has a unique maximum

⁸Section 3 focuses on normal distributed ξ_{it} in Examples 1 & 2 satisfying the completeness assumptions with two type of DGPs, whereas subsection 4.3 presents a test concerning the completeness of $f_{Y_{it}|X_{it},Y_{it-1},X_{it-1}}$ in Assumptions 2.6.

at the true parameter θ_0 . In a parametric likelihood case,⁹ the local identifiability of the unknown parameter vector is equivalent to the non-singularity of the information matrix under weak regularity conditions. If the true parameter θ_0 is a critical point of $K(\theta_0)$, then a sufficient condition of the uniqueness of θ_0 is $K''(\theta_0)$ is negative semidefinite. The second derivative of $K(\theta_0)$ in the scalar case is

$$K''(\theta_0) = -E \left[\left(\frac{\frac{\partial}{\partial \theta} f_{X_{it}, Y_{it-1}, X_{it-1}; \theta} |_{\theta=\theta_0}}{f_{X_{it}, Y_{it-1}, X_{it-1}}} \right)^2 \right] \quad (21)$$

where the expression of $\frac{\partial}{\partial \theta} f_{X_{it}, Y_{it-1}, X_{it-1}; \theta} |_{\theta=\theta_0}$ appears in Eq. (67). The vector case of $K''(\theta_0)$ is $K''(\theta_0) = [K_{lm}]$, where

$$K_{lm} = -E \left[\left(\frac{\frac{\partial}{\partial \theta_l} f_{X_{it}, Y_{it-1}, X_{it-1}; \theta} |_{\theta=\theta_0} \frac{\partial}{\partial \theta_m} f_{X_{it}, Y_{it-1}, X_{it-1}; \theta} |_{\theta=\theta_0}}{f_{X_{it}, Y_{it-1}, X_{it-1}}^2} \right) \right], \quad (22)$$

where $\frac{\partial}{\partial \theta_l} f_{X_{it}, Y_{it-1}, X_{it-1}; \theta} |_{\theta=\theta_0}$ is equal to the term in Eq. (67) after replacing with the partial derivative $\frac{\partial}{\partial \theta_l}$. These results are sufficient conditions for the identification.

Corollary 2.2. *Suppose that in an open neighbor of θ_0 in Θ , the second derivative of the Kullback-Leibler function $K(\theta)$ in Eq. (21) or (22) is negative definite. Under Assumptions 2.3, 2.4(ii) and 2.5-2.6, the semi-parametric censored density function $f_{Y_{it}|X_{it}, Y_{it-1}, U_{it}; \theta}$ and the joint density function $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}}$ can be identified given the distribution of the two-period observable variables $(y_{it}, x_{it}, y_{it-1}, x_{it-1})$ for $i = 1, 2, \dots, N$.*

Proof: See the appendix.

3. Examples

Consider the two examples presented at the beginning of Section 2. This section shows when the completeness conditions in Section 2 hold in these cases. Assumptions 2.3, 2.4(ii), and 2.5 are related to the completeness of the variant forms of the semi-parametric censored density function $f_{Y_{it}^+|X_{it}, Y_{it-1}, U_{it}; \theta}$. Equations (7) and (9) show that the completeness of the semi-parametric censored density functions over positive Y_{it} in the two motivating examples are connected to the PDF of the random shock ξ_{it} . Therefore, this section focuses on what kind of

⁹The parameter θ only contains a finite-dimensional component.

semi-parametric distribution assumptions in ξ_{it} make these examples satisfy the completeness assumptions. For simplicity, assume the domains of ξ_{it} and U_{it} are \mathbb{R} .

Most of the interesting leading cases for Models (7) and (9) occur when the random shock ξ_{it} is assumed to have an independent Gaussian white noise process.¹⁰ For simplicity, assume $g_2(X'_{it}, Y_{it-1}; \theta_2) = X'_{it}\beta + \gamma Y_{it-1}$. In this case, the semi-parametric censored density function $f_{Y_{it}|X_{it}, Y_{it-1}, U_{it}}$ is fully parameterized and correctly specified at θ_0 . The specifications of the models under the normality assumption are as follows:

Semi-parametric Dynamic Tobit Models:

Assuming that $\xi_{it} \sim N(0, \sigma_\xi)$, Eq. (7) leads to

$$f_{Y_{it}|X_{it}, Y_{it-1}, U_{it}; \theta} = \left[1 - \Phi \left(\frac{X'_{it}\beta + \gamma Y_{it-1} + U_{it}}{\sigma_\xi} \right) \right]^{\mathbf{1}(Y_{it}=0)} \times \left[\frac{1}{\sigma_\xi} \phi \left(\frac{Y_{it} - X'_{it}\beta - \gamma Y_{it-1} - U_{it}}{\sigma_\xi} \right) \right]^{\mathbf{1}(Y_{it}>0)}, \quad (23)$$

where $\theta = (\beta, \gamma, \sigma_\xi^2)^T$.

Semi-parametric Dynamic Lognormal Hurdle Models:

Let $g_3(X'_{it}, Y_{it-1}; \theta_1) = X'_{it}\beta_d + \gamma_d Y_{it-1}$. Suppose that $\varsigma_{it} \sim N(0, 1)$ and $\xi_{it} \sim N(0, \sigma_\xi)$. Equation (9) then becomes

$$f_{Y_{it}|X_{it}, Y_{it-1}, U_{it}; \theta} = (1 - \Phi(X'_{it}\beta_d + \gamma_d Y_{it-1} + U_{it}))^{\mathbf{1}(Y_{it}=0)} \left\{ \Phi(X'_{it}\beta_d + \gamma_d Y_{it-1} + U_{it}) \times \phi \left(\frac{\log(Y_{it}) - X'_{it}\beta - \gamma Y_{it-1} - U_{it}}{\sigma_\xi} \right) \frac{1}{\sigma_\xi Y_{it}} \right\}^{\mathbf{1}(Y_{it}>0)}, \quad (24)$$

where $\theta = (\beta_d, \gamma_d, \beta, \gamma, \sigma_\xi)^T$.

The normality assumption makes it possible to verify the completeness of the semi-parametric censored density function $f_{Y_{it}^+|X_{it}, Y_{it-1}, U_{it}; \theta}$ in Assumptions 2.3, 2.4(ii), and 2.5 directly. It is then necessary to show that the semi-parametric censored density functions (23) and (24) satisfy these completeness assumptions. To do this, introduce the completeness of normal distributions and exponential families in \mathcal{L}^2 from Hu and Shiu (2011b) which are

¹⁰There may exist more different types of the distributions for the random shock ξ_{it} and the normality assumption here only illustrates the application of the results in 2.1.

variants of the results of Newey and Powell (2003).¹¹

Lemma 3.1. *Suppose that the distribution of u conditional on y is $N(a+by, \sigma^2)$ for $b, \sigma^2 > 0$ and the support of y contains an open set. In this case, $E[h(\cdot)|y] = 0$ for any $x \in \mathcal{Y}$ implies $h(u) = 0$ almost everywhere in \mathcal{U} ; equivalently, $\{f(u|y) : y \in \mathcal{Y}\}$ is complete in $\mathcal{L}^2(\mathcal{U})$.*

Lemma 3.2. *Let $f(u|y) = s(u)t(y) \exp[\mu(y)\tau(u)]$, where $s(u) > 0$, $\tau(u)$ is one-to-one in u , and the support of $\mu(y)$, \mathcal{Y} , contains an open set. In this case, $E[h(\cdot)|y] = 0$ for any $y \in \mathcal{Y}$ implies $h(u) = 0$ almost everywhere in \mathcal{U} ; equivalently, the family of conditional density functions $\{f(u|y) : y \in \mathcal{Y}\}$ is complete in $L^2(\mathcal{U})$.*

Lemma 3.1 implies that for an open set $O_y \subset \mathcal{Y}$, $\{\phi\left(\frac{u-(a+by)}{\sigma}\right) : y \in O_y\}$ is complete in $\mathcal{L}^2(\mathcal{U})$. This completeness can be extended to a weighted space $\mathcal{L}^2(\mathcal{U}, \omega)$ for an appropriately chosen ω . Set $\omega(u) = e^{-\frac{u^2}{2\sigma^2}}$. Suppose that $h \in \mathcal{L}^2(\mathcal{U}, \omega)$ such that for $y \in O_y$,

$$\int h(u)\phi\left(\frac{u-(a+by)}{\sigma}\right)\omega(u)du = 0.$$

Multiplying the equation by $e^{\frac{1}{4\sigma^2}(a+by)^2}$ results in

$$0 = \int h(u)\phi\left(\frac{u-(a+by)}{\sigma}\right)\omega(u)e^{\frac{1}{4\sigma^2}(a+by)^2}du.$$

It follows that

$$0 = \int h(u)\omega(u)^{\frac{1}{2}}\phi\left(\frac{u-\frac{1}{2}(a+by)}{\sigma/\sqrt{2}}\right)du$$

for $y \in O_y$. Note $h(u)\omega(u)^{\frac{1}{2}} \in \mathcal{L}^2(\mathcal{U})$ because $h \in \mathcal{L}^2(\mathcal{U}, \omega)$. Lemma 3.1 also implies that $\{\phi\left(\frac{u-\frac{1}{2}(a+by)}{\sigma/\sqrt{2}}\right) : y \in O_y\}$ is complete $L^2(\mathcal{U})$. Applying this result to the equation suggests that $h(u) = 0$ almost everywhere. Therefore, $\{\phi\left(\frac{u-(a+by)}{\sigma}\right) : y \in O_y\}$ is complete in $L^2(\mathcal{U}, \omega)$.

Based on the information about the completeness of normal distributions, it is possible to investigate the completeness condition of Models (23) and (24).

Semi-parametric Dynamic Tobit Models:

¹¹See Theorems 2.2 and 2.3 in Newey and Powell (2003) for details. More general discussions of completeness can be found in D'Haultfoeuille (2011), Andrews (2011), and Hu and Shiu (2011b).

Set $\tilde{\mathcal{Y}}_{it} = \mathbb{R}^+$. Given $\theta \in \Theta$, and (x_{it}, y_{it-1}) ,

$$f_{Y_{it}^+|X_{it}, Y_{it-1}, U_{it}; \theta} = \frac{1}{\sigma_\xi} \phi \left(\frac{Y_{it}^+ - X_{it}'\beta - \gamma Y_{it-1} - U_{it}}{\sigma_\xi} \right). \quad (25)$$

Semi-parametric Dynamic Lognormal Hurdle Models:

Given $\theta \in \Theta$, and (x_{it}, y_{it-1}) ,

$$f_{Y_{it}^+|X_{it}, Y_{it-1}, U_{it}; \theta} = \Phi \left(X_{it}'\beta_d + \gamma_d Y_{it-1} + U_{it} \right) \phi \left(\frac{\log(Y_{it}^+) - X_{it}'\beta - \gamma Y_{it-1} - U_{it}}{\sigma_\xi} \right) \frac{1}{\sigma_\xi Y_{it}^+} \quad (26)$$

The completeness conditions in Section 2 are all associated with the dependent variables Y_{it}^+ and the unobserved covariate U_{it} . Therefore, it is necessary to investigate which functional forms connect these two variables. In these models, the dependent variables Y_{it}^+ and the unobserved covariate U_{it} are both inside the standard normal PDF ϕ . It follows that semi-parametric dynamic tobit models satisfies Assumptions 2.3 and 2.4(ii) by switching the role between Y_{it}^+ and U_{it} to the result of Lemma 3.1. Because the standard normal CDF Φ is positive, the semi-parametric dynamic lognormal hurdle models also fulfill Assumptions 2.3 and 2.4(ii) using Lemma 3.1 and a change of variable.¹²

Assumption 2.5 requires that the partial derivatives of the semi-parametric censor density function with respect to all components of the parameter θ be complete. According to the functional forms in Eqs. (25) and (26) and use of a change of variable, two types of the partial derivatives of $f_{Y_{it}^+|X_{it}, Y_{it-1}, U_{it}; \theta_0}$ should be considered. The first one is the partial derivative with respect to the components of β and γ , and the second one is σ_ξ . The completeness of the first type can be reduced to the completeness of the family of $\{(y - c - u) \phi \left(\frac{y - c - u}{\sigma_\xi} \right) : u \in \mathcal{U}\}$ in $L^2(\tilde{\mathcal{Y}})$ for some constant c . Similarly, the completeness of the second type depends on the family of $\{(\sigma_\xi^2 - (y - c - u)^2) \phi \left(\frac{y - c - u}{\sigma_\xi} \right) : u \in \mathcal{U}\}$ in $L^2(\tilde{\mathcal{Y}})$ for some constant c . The following lemma provides the completeness of the families of variant of the normal PDF ϕ .

Lemma 3.3. *Suppose the domain \mathcal{U} contains an open set. For a constant c , the families of functions $\{(y - c - u) \phi \left(\frac{y - c - u}{\sigma_\xi} \right) : u \in \mathcal{U}\}$ and $\{(\sigma_\xi^2 - (y - c - u)^2) \phi \left(\frac{y - c - u}{\sigma_\xi} \right) : u \in \mathcal{U}\}$ are*

¹²Use $\dot{Y}_{it} = \log(Y_{it}^+)$ and then $d\dot{Y}_{it} = \frac{1}{Y_{it}^+} dY_{it}^+$.

complete in $L^2(\tilde{\mathcal{Y}})$.

Proof: See the appendix.

This discussion also applies to models with heteroskedasticity, which allow more general functional form in corner solution models. If ξ_{it} has a heteroskedastic normal distribution such that $\xi_{it} \sim N(0, h(X'_{it}, Y_{it-1}; \sigma_\xi))$ then the semi-parametric censored density functions in Eqs. (23) and (24) respectively become

$$f_{Y_{it}|X_{it}, Y_{it-1}, U_{it}; \theta} = \left[1 - \Phi \left(\frac{X'_{it}\beta + \gamma Y_{it-1} + U_{it}}{h(X'_{it}, Y_{it-1}; \sigma_\xi)} \right) \right]^{\mathbf{1}(Y_{it}=0)} \times \left[\frac{1}{h(X'_{it}, Y_{it-1}; \sigma_\xi)} \phi \left(\frac{Y_{it} - X'_{it}\beta - \gamma Y_{it-1} - U_{it}}{h(X'_{it}, Y_{it-1}; \sigma_\xi)} \right) \right]^{\mathbf{1}(Y_{it}>0)}, \quad (27)$$

and

$$\begin{aligned} f_{Y_{it}|X_{it}, Y_{it-1}, U_{it}; \theta} &= (1 - \Phi(X'_{it}\beta_d + \gamma_d Y_{it-1} + U_{it}))^{\mathbf{1}(Y_{it}=0)} \left\{ \Phi(X'_{it}\beta_d + \gamma_d Y_{it-1} + U_{it}) \right. \\ &\quad \times \left. \phi \left(\frac{\log(Y_{it}) - X'_{it}\beta - \gamma Y_{it-1} - U_{it}}{h(X'_{it}, Y_{it-1}; \sigma_\xi)} \right) \frac{1}{h(X'_{it}, Y_{it-1}; \sigma_\xi) Y_{it}} \right\}^{\mathbf{1}(Y_{it}>0)}. \end{aligned} \quad (28)$$

Adding the heterogeneous structure does not affect the functional form, which dominates both the dependent variables Y_{it}^+ and the unobserved covariate U_{it} . The derivations in homoskedastic cases can be extended to heteroskedastic cases without difficulty.

The assumptions not related to the semi-parametric censored density functions include Assumptions 2.4(i) and 2.6. These assumptions require functional form restrictions on the conditional density function $f_{Y_{it}^+|X_{it}, Y_{it-1}, X_{it-1}}$ of observables. With the well-known completeness from the normal distributions and the exponential families in Lemma 3.1 and Lemma 3.2, it is possible to construct two types of $f_{Y_{it}^+|X_{it}, Y_{it-1}, X_{it-1}}$ satisfying Assumptions 2.4(i) and 2.6. Assumption 2.6 implies Assumption 2.4(i). Given a fixed (X_{it}, Y_{it-1}) , suppose \mathcal{X}_{it-1} contains an open set. If $f_{Y_{it}^+|X_{it}, Y_{it-1}, X_{it-1}} = \phi(Y_{it}^+ - \psi_1(X_{it}, Y_{it-1}; \theta_1) - \beta\psi_2(X_{it-1}))$, then

$$\begin{aligned} \frac{\partial}{\partial X_{it-1}} f_{Y_{it}^+|X_{it}, Y_{it-1}, X_{it-1}} &= \beta\psi'_2(X_{it-1}) (Y_{it}^+ - \psi_1(X_{it}, Y_{it-1}; \theta_1) - \beta\psi_2(X_{it-1})) \phi(Y_{it}^+ - \psi_1(X_{it}, Y_{it-1}; \theta_1) - \beta\psi_2(X_{it-1})). \end{aligned} \quad (29)$$

A sufficient condition to satisfy Assumption 2.6 for this specification of $f_{Y_{it}^+|X_{it},Y_{it-1},X_{it-1}}$ is $\beta\psi_2'(X_{it-1}) \neq 0$ and the range of ψ_2 contains an open set, according to Lemma 3.3. On the other hand, consider

$$\begin{aligned} & f_{Y_{it}^+|X_{it},Y_{it-1},X_{it-1}} \\ &= s(Y_{it}^+, X_{it}, Y_{it-1})t(X_{it}, Y_{it-1}, X_{it-1}) \exp [\mu(X_{it}, Y_{it-1}, X_{it-1})\tau(Y_{it}^+, X_{it}, Y_{it-1})] \end{aligned}$$

where $s(Y_{it}^+, X_{it}, Y_{it-1}) > 0$, $\tau(Y_{it}^+, X_{it}, Y_{it-1}) \neq 0$ is one-to-one in Y_{it}^+ , and the support of μ contains an open set such that $\frac{\partial}{\partial X_{it-1}}\mu(X_{it}, Y_{it-1}, X_{it-1}) \neq 0$. This conditional density function also fulfills Assumption 2.4(i) and Assumption 2.6.

The examples in this section rely on the normality of the random shock ξ_{it} and it is possible to relax the normality assumption. However, in limited dependent variable models, the key issue is comparing estimated average partial effects across different models rather than parameter estimates. These models are likely to do an appropriate job of providing average partial effects under more general settings.

4. Semiparametric Estimation and Inference

The semi-parametric censored density function (10) identified in Theorem 2.1 can be determined using Eq. (12). Optimizing certain empirical criteria in general parameter spaces produces a sieve maximum likelihood estimator (sieve MLE). The integral Eq. (12) suggests a corresponding sieve MLE:

$$\begin{aligned} & (\hat{\theta}, \hat{f}_1)^T \\ &= \arg \max_{(\theta, f_1)^T \in \mathcal{A}_n} \frac{1}{N} \sum_{i=1}^N \ln \int f_{Y_{it}|X_{it},Y_{it-1},U_{it};\theta}(y_{it}|x_{it}, y_{it-1}, u_{it}) f_1(x_{it}, y_{it-1}, x_{it-1}, u_{it}; \theta) du_{it}, \end{aligned} \tag{30}$$

using a two-period i.i.d. sample $\{y_{it}, x_{it}, y_{it-1}, x_{it-1}\}_{i=1}^N$.¹³ The space \mathcal{A}_n is a sequence of approximating sieve spaces containing sieve approximations of the parameter because maximization over the whole parameter space \mathcal{A} is undesirable. In addition, θ is a finite-dimensional parameter of interest and f_1 is a potentially infinite-dimensional nuisance parameter or non-

¹³A general review of semi-parametric sieve MLE appears in Shen (1997), Chen and Shen (1998), and Ai and Chen (2003).

parametric component that varies with θ . The following subsection provides a detailed implementation of sieve approximations of the nonparametric component f_1 .

4.1. Restrictions on Sieve Coefficients

As for a nonparametric series estimator of $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta}$, constructing a sieve approximating series that varies with the model parameter θ is an essential issue for the proposed sieve MLE. The sieve expression of $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \delta_1}$ in dynamic censored models with a lagged dependent variable consists of two different parts, $Y_{it-1} = 0$ and $Y_{it-1} > 0$, and these parts can be build according to their numerical structures. Set $f_{Y_{it-1}=0} = \text{Prob}(Y_{it-1} = 0)$. A way to split these two parts is

$$f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \delta_1} = \begin{cases} f_{X_{it}, X_{it-1}, U_{it} | Y_{it-1}=0} f_{Y_{it-1}=0} & \text{if } y = 0, \\ f_{X_{it}, Y_{it-1}>0, X_{it-1}, U_{it}} & \text{if } y > 0. \end{cases}$$

The corresponding density restrictions are

$$\int f_{X_{it}, X_{it-1}, U_{it} | Y_{it-1}=0} dx_{it} dx_{it-1} du_{it} = 1, \text{ and} \\ f_{Y_{it-1}=0} + \int f_{X_{it}, Y_{it-1}>0, X_{it-1}, U_{it}} dy_{it-1} dx_{it} dx_{it-1} du_{it} = 1.$$

Set $z_{1, \sigma_\xi} \equiv \frac{x'_{it}\beta - x'_{it-1}\beta - u_{it}}{\sigma_\xi}$, and $z_{2, \sigma_\xi} \equiv \frac{x'_{it}\beta - u_{it}}{\sigma_\xi}$. For the $Y_{it-1} = 0$ part, consider

$$(f_{X_{it}, X_{it-1}, U_{it} | Y_{it-1}=0})^{1/2} = \sum_{i=0}^{i_n} \sum_{j=0}^{j_n} \sum_{k=0}^{k_n} \hat{a}_{ijk} q_i(z_{1, \sigma_\xi}) q_j(z_{2, \sigma_\xi}) q_k\left(\frac{u_{it}}{\sigma_\xi}\right).$$

where q'_i 's, q'_j 's, and q'_k 's represent the orthonormal Fourier series:

$$q_0(z_1) = \frac{1}{\sqrt{l_1}} \text{ and } q_i(z_1) = \frac{1}{\sqrt{l_1}} \sin\left(\frac{i\pi}{l_1} z_1\right) \text{ or } q_i(z_1) = \frac{1}{\sqrt{l_1}} \cos\left(\frac{i\pi}{l_1} z_1\right), \\ q_0(z_2) = \frac{1}{\sqrt{l_2}} \text{ and } q_j(z_2) = \frac{1}{\sqrt{l_2}} \sin\left(\frac{j\pi}{l_2} z_2\right) \text{ or } q_j(z_2) = \frac{1}{\sqrt{l_2}} \cos\left(\frac{j\pi}{l_2} z_2\right), \\ q_0(u_{it}) = \frac{1}{\sqrt{l_3}}, q_k(u_{it}) = \sqrt{\frac{2}{l_3}} \cos\left(\frac{k\pi}{l_3} u_{it}\right),$$

On the other hand, suppose that $y_{it-1} \in (0, l_4]$. Write

$$(f_{X_{it}, Y_{it-1} > 0, X_{it-1}, U_{it}})^{1/2} = \sum_{i=0}^{i_n} \sum_{j=0}^{j_n} \sum_{k=0}^{k_n} \sum_{l=0}^{l_n} \tilde{a}_{ijkl} \tilde{q}_i(z'_{1, \sigma_\xi}) \tilde{q}_j(z'_{2, \sigma_\xi}) \tilde{q}_k\left(\frac{u_{it}}{\sigma_\xi}\right) \tilde{q}_l\left(\frac{y_{it-1}}{l_4}\right),$$

where $z'_{1, \sigma_\xi} \equiv \frac{x'_{it}\beta - \gamma y_{it-1} - x'_{it-1}\beta - u_{it}}{\sigma_\xi}$, $z'_{2, \sigma_\xi} \equiv \frac{x'_{it}\beta - \gamma y_{it-1} - u_{it}}{\sigma_\xi}$, and $q_0(z_4) = \frac{1}{\sqrt{l_4}}$, $q_l(z_4) = \sqrt{\frac{2}{l_4}} \cos(\frac{l\pi}{l_4} z_4)$.

The density restrictions for these sieve coefficients are

$$\sum_{i=0}^{i_n} \sum_{j=0}^{j_n} \sum_{k=0}^{k_n} (\hat{a}_{ijk})^2 = 1 \text{ and } f_{Y_{it-1}=0} + \sum_{i=0}^{i_n} \sum_{j=0}^{j_n} \sum_{k=0}^{k_n} \sum_{l=0}^{l_n} (\tilde{a}_{ijkl})^2 = 1. \quad (31)$$

4.2. Estimating Average Partial Effects

Denote $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \hat{\theta}, \delta_1}$ as the sieve MLE of the initial joint distribution $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}}$ in the dynamic tobit model, where $\hat{\theta}$ is the estimated finite dimensional parameter of the proposed sieve MLE. This parameter can be used to obtain the sieve approximations of the marginal distribution of the unobserved covariate U_{it} :

$$\hat{f}_{U_{it}} = \int_{\mathcal{X}_{it}} \int_{\mathcal{Y}_{it-1}} \int_{\mathcal{X}_{it-1}} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \hat{\theta}, \delta_1} dx_{it} dy_{it-1} dx_{it-1} \quad (32)$$

Therefore, under the assumptions made in Theorem 2.1, it is possible to consistently estimate average partial effects at interesting values of the explanatory variables. The average structural functions in the dynamic tobit models are based on

$$\begin{aligned} \hat{\mu}(X_{it}, Y_{it-1}) &\equiv \int_{U_{it}} \left(\int_{Y_{it}} \max(0, y_{it}) f_{Y_{it}|X_{it}, Y_{it-1}, U_{it}; \hat{\theta}} dY_{it} \right) \hat{f}_{1, U_{it}} dU_{it} \\ &= \int_{U_{it}} \left[\Phi \left(\frac{X'_{it}\hat{\beta} + \hat{\gamma}Y_{it-1} + U_{it}}{\hat{\sigma}_\xi} \right) (X'_{it}\hat{\beta} + \hat{\gamma}Y_{it-1} + U_{it}) \right. \\ &\quad \left. + \hat{\sigma}_\xi \phi \left(\frac{X'_{it}\hat{\beta} + \hat{\gamma}Y_{it-1} + U_{it}}{\hat{\sigma}_\xi} \right) \right] \hat{f}_{1, U_{it}} dU_{it}. \end{aligned} \quad (33)$$

The magnitude of state dependence or average partial effect from $Y_0 = 0$ to Y_1 at interesting values of the explanatory variable X_{it} can be measured by the difference

$$\hat{\mu}(X_{it}, Y_1) - \hat{\mu}(X_{it}, Y_0 = 0). \quad (34)$$

On the other hand, the average partial effect (APE) of a continuous explanatory variable can be defined using derivatives of the average structural functions in Eq. (33).

4.3. Inference

This study addresses two inference problems. One is to provide standard errors of the estimated finite dimensional parameter $\hat{\theta}$ and the other one is to test the completeness of Assumptions 2.4(i) and 2.6. The parameter $\hat{\theta}$ is estimated by optimization methods for the integrated nonlinear objective function in Eq. (30). Obtaining a formula of standard errors for this sieve MLE estimator is somewhat complicated, but it can be done using bootstrapping. Chen, Linton, and Van Keilegom (2003) studies sufficient conditions for the consistency and asymptotic normality of a class of semiparametric optimization estimators in which the criterion function does not obey the standard smoothness condition. Their results prove the validity of the bootstrap method for estimating correct confidence regions for the finite dimensional parameter θ asymptotically.

On the other hand, because the key conditional distribution $f_{Y_{it}^+|X_{it},Y_{it-1},X_{it-1}}$ in Assumptions 2.4(i) and 2.6 is observable, the conditions seem to be directly testable using both two periods of data. However, in most cases, the exact form of the conditional distribution is not obvious and it is difficult to test nonparametrically with continuous variables.¹⁴ To provide more support for the validity of the condition, this study presents a test based on a parametric setting and uses the data to estimate unknown parameters associated with the completeness under parametric specifications. Although there are many ways to model completeness, this study adopts a fairly flexible approach. The previous section provides an example of completeness in which $f_{Y_{it}^+|X_{it},Y_{it-1},X_{it-1}}$ is normally distributed. Given functions ψ_1 and ψ_2 such that $\psi_2' \neq 0$, suppose $\psi_2(\mathcal{X}_{it-1})$ contains an open set. Consider

$$f_{Y_{it}^+|X_{it},Y_{it-1},X_{it-1}} = \phi(Y_{it}^+ - \psi_1(X_{it}, Y_{it-1}; \theta_1) - \beta\psi_2(X_{it-1})).$$

As shown in Eq. (29), under the parametric representation, the completeness in both Assumptions 2.4(i) and 2.6 depends on whether the parameter $\beta\psi_2'(X_{it-1})$ is equal to zero. The

¹⁴See Canay, Santos, and Shaikh (2011) for details.

null hypothesis that $f_{Y_{it}^+|X_{it},Y_{it-1},X_{it-1}}$ fails Assumptions 2.4(i) and 2.6 is

$$H_0 : \beta = 0.$$

When X_{it}, X_{it-1} are vectors of covariates, it is possible to conduct these tests on continuously distributed covariates. If we model $\psi_1(X_{it}, Y_{it-1}; \theta_1)$ as a linear function of the regressors, a standard t statistics to test H_0 can be conducted using OLS regression.

5. Monte Carlo Simulation

This study presents the finite sample property of the proposed sieve MLE estimators based on a Monte Carlo simulation study. The simulation design in this study is similar to the dynamic panel data models discussed by Shiu and Hu (2010). The proposed model adopts the following procedure:

Dynamic tobit Models with AR(1) Transitory Error:

$$Y_{it} = \max\{\beta_0 + \beta_1 X_{it} + \gamma Y_{it-1} + V_i + \varepsilon_{it}, 0\} \quad i = 1, \dots, N; t = 1, \dots, T - 1. \quad (35)$$

where $V_i \sim N(1, 1/2)$ and $\varepsilon_{it} = \rho \varepsilon_{it-1} + \xi_{it}$ with $\xi_{it} \sim N(0, \sigma_\xi^2)$. The unobserved covariate $U_{it} = V_i + \rho \varepsilon_{it-1}$. As discussed earlier, the models can be transformed into $f_{Y_{it}|X_{it},Y_{it-1},U_{it};\theta}$ by Eq. (23). Set $h(x) = 0.2 \exp(-x)$. The generating processes of the covariate evolution have the following form $X_{it+1} = X_{it} + h(X_{it})\eta_{it} + U_{it}$ with $\eta_{it} \sim N(0, 1)$ or

$$f_{X_{it+1}|X_{it},U_{it}}(x_{it+1}|x_{it},u_{it}) = \frac{1}{h(x_{it})} \phi_\eta \left(\frac{x_{it+1} - x_{it} - u_{it}}{h(x_{it})} \right),$$

where ϕ_η is the standard normal distribution. Four different data generating processes (DGP) are as follows:

$$\begin{aligned} \text{DGP I:} \quad & (\beta_0, \beta_1, \gamma, \sigma_\xi^2, \rho) = (0.2, -1, 0, 0.5, 0) \\ \text{DGP II:} \quad & (\beta_0, \beta_1, \gamma, \sigma_\xi^2, \rho) = (0.2, -1, 0, 0.5, 0.5) \\ \text{DGP III:} \quad & (\beta_0, \beta_1, \gamma, \sigma_\xi^2, \rho) = (0.2, -1, 1, 0.5, 0.5) \\ \text{DGP IV:} \quad & (\beta_0, \beta_1, \gamma, \sigma_\xi^2, \rho) = (0.2, -1, 1, 0.5, -0.5). \end{aligned}$$

In the all designs, set $\beta_0 = 0.2$, $\beta_1 = -1$ and $\sigma_\xi^2 = 0.5$. These designs focus on different values for state dependence, γ , and AR(1) coefficient of the serially correlated error term, ρ . The simulation designs in DGPs I & II do not have state dependence ($\gamma = 0$), but the simulation designs in DGPs III & IV show strong persistent effects from the past dependent variable ($\gamma = 1$). This study assumes that the panel data is set in three different observations, 250, 500, 1000, and presents experiments for small T for which the sampling data are drawn over $T = 3$ periods.

The two main differences between this experiment and the study by Shiu and Hu (2010) is that 1) at least three periods of data are needed for the simulation in Shiu and Hu (2010), and 2) the generating processes of the covariate evolution $f_{X_{it+1}|X_{it},U_{it}}$ is required to satisfy a mode condition, which is one of the nonparametric identification assumptions in Shiu and Hu (2010). Thus, the current estimation results cannot be compared to the results of Shiu and Hu (2010) if we only use a two-period simulated sample. Another practical advantage of the proposed sieve MLE estimator is that it does not require sieve implementation of the covariate evolution. Hence, the implementation is easier but the normality assumption of ξ_{it} is required and essential to the estimation. The proposed method uses the Fourier series in Subsection 4.2 with the number of term, $i_n = 5$, $j_n = 2$, $k_n = 2$, and $l_n = 2$, to approximate the initial joint distribution $f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta}$. Two-period or three-period simulated samples are used to conduct estimations for 100 replications.

Tables 1-3 present simulation results for the model parameters of the dynamic tobit model. These tables present the means and the medians in estimating β_0 , β_1 , σ_ξ^2 , and γ together with their standard errors. In calculating the standard error of coefficient estimators, use the variance of coefficients estimated from the 100 replications as a measure of true variance. Observe the following patterns in the simulation results for the coefficient estimators. First, there generally exists downward bias in estimating the autoregressive parameter γ in all sample sizes. Second, the bias in estimating all parameters ($\beta_0, \beta_1, \sigma_\xi^2$) is small, suggesting that the proposed sieve MLE estimator achieves consistent estimation results. Finally, all standard errors do not vary much. This suggests that for these samples, the statistical performances of the proposed sieve MLE estimator are very close. This study presents a comparison of the proposed sieve MLE estimators with the benchmark estimator and the three-period estimator of Shiu and Hu (2010). The benchmark estimator treats the unobserved U_{it} as a covariate

and applies a MLE method. As expected, this benchmark estimator performs better than the other two estimators, and the biases and standard errors decrease in larger samples. In DPGs with nontrivial state dependence, the three-period estimators provides better estimators for γ . This suggests that in the same sample sizes, an additional period of data may help reveal the dynamic structure of the data.

Table 4 presents the magnitude of the state dependence $SD(\bar{X}_{it})$. The results in DGP I & II imply that the proposed sieve MLE performs well because the parameter $\gamma = 0$ in these cases and the estimation results are close to zero. The DGP with nontrivial positive state dependence exhibit significant variation in the estimation results of $SD(\bar{X}_{it})$, and the average response of the state dependence for these DGPs is at least approximately 0.5. In addition, the means and medians of $SD(\bar{X}_{it})$ are similar, reflecting some symmetry in their respective distributions.

6. Empirical Application

This study reports the application of the proposed sieve MLE estimator to a censored dynamic tobit model describing the annual health expenditures of individuals given their past health expenditures and other covariates. In this case, the dependent variables represent the log values of annual individual medical expenditures plus one. To accommodate the piles of the corner outcomes, this censored dynamic tobit model is a natural fit for this health expenditure topic. Identification results show that the proposed model has some advantages: (i) arbitrary correlation between unobserved time invariant factors, such as individual inherent health and other explanatory variables, and (ii) allowing the absence of initial observations of individual health expenditures. In addition, the proposed sieve MLE estimator only requires two periods of data and provides average partial effects.

The empirical analysis in this study is based on the Medical Expenditure Panel Survey (MEPS) Panel 4. The MEPS data provides nationally representative information on health care use, expenditures, sources of payment, and insurance coverage for the U.S. population from 1,999 to 2,000. The MEPS, which contains detailed data on annual total health care expenditure, demographic characteristics, health conditions, health status, use of medical care services, and income, is appropriate for our empirical application. Table 5 presents

summary statistics of health insurance variables, socioeconomic variables, and health status regressors for the first-year and the second-year of the data. We have a two-periods of the data with 7,669 cross-sectional observations. There are sizable fractions of the sample with zero medical expenditure, 18.646% (1,430/7,669) and 20.576% (1,578/7,669) in Periods 1 and 2, respectively.

The estimated equation of dynamic health expenditures is

$$\begin{aligned} Y_{it} &= \max \left\{ 0, X'_{it}\beta + \gamma Y_{it-1} + V_i + \underbrace{\eta_{it} + \xi_{it}}_{\varepsilon_{it}} \right\} \quad \forall i = 1, \dots, N; t = 1, 2, \\ &= \max \left\{ 0, X'_{it}\beta + \gamma Y_{it-1} + U_{it} + \xi_{it} \right\} \end{aligned} \quad (36)$$

The dependent variable $Y_{it} = \text{Lnexp}_{it}$ is the natural logarithm of health expenditure plus one. The covariate $X_{it} = \left(\text{Lninc}_{it}, \text{Lnfam}_{it}, \text{Age}_{it}, \text{Male}_{it}, \text{Black}_{it}, \text{Education}_{it}, \text{Phsical}_{it}, \text{Ndental}_{it}, \text{Good}_{it}, \text{Fair}_{it}, \text{Poor}_{it}, \dots, \text{Time dummies} \right)$. The unobserved heterogeneity V_i represents time-invariant individual heterogeneity factors, such as inherent ability or personal regimen to resist negative health shock. Assume that Assumptions 2.1 and 2.2 split the transitory error term ε_{it} into η_{it} and ξ_{it} , and that ξ_{it} is normally distributed. This normality assumption guarantees that Assumptions 2.3, 2.4(ii), and 2.5 are fulfilled, as Section 3 shows. Assumptions 2.4(i) and 2.6 demand the completeness conditions related to the family of conditional distribution of positive health expenditure y_{it}^+ over x_{it-1} , $\{f_{Y_{it}^+|X_{it}, Y_{it-1}, X_{it-1}} : x_{it-1} \in \mathcal{X}_{it-1}\}$. Choose ψ_1 as a linear function and ψ_2 as a linear function of squares to conduct the testing proposed in Subsection 4.3. The estimated coefficient of the squared Lninc_{it-1} is -0.0014 with a p -value of 0.052. In an intuitive sense, the testing result suggests that the covariate at the previous period x_{it-1} containing income squares has enough variation such that the conditional distribution of positive health expenditure can cover all variation of positive health expenditures. These assumptions, along with the mild regularity condition stated in Theorem 2.1, provide the identification of Model (36) and the sieve MLE developed in Section 4 is applicable.

Table 6 shows the results of the estimation of panel data Model (36) using three specifications, including a static linear fixed effect model (Column 1), a static tobit model with random effect (Column 2), and the semiparametric dynamic tobit model (Column 3). The

three sets of estimates present similar results in terms of directions of effects and estimated coefficients. As expected, there are differences in the magnitudes of the estimated APEs in the RE tobit and semiparametric dynamic tobit specifications. The APEs of semiparametric dynamic tobit specifications have greater effects after controlling for the dynamic effect of health expenditures. The coefficient estimates of state dependence effect of health expenditures is up to 1.052. As a result, the effect of previous health expenditures on the future health expenditures is estimated to be, in APE, 1.448. The estimated coefficient shows that the previous health expenditures have persistent effects or there is large first order state dependence of health expenditures. One of the variables of interest is $Lninc_{it}$, the natural logarithm of the family income plus one. The coefficient of $Lninc_{it}$ in regression on $Lninc_{it}$ represents the income elasticity of demand for health care. The result of the semiparametric dynamic tobit specifications indicates that individuals consume more health care when their incomes go up after controlling for the past health expenditures.

7. Conclusion

This study presents identification results for the semi-parametric censored dynamic panel data models and their corresponding average partial effects. The main assumptions of the proposed method include the existence of an independent random shock, a semi-parametric specification of the random shock, and the completeness of families of known PDFs corresponding to censored dynamic panel data models and observed conditional density functions between the dependent and explanatory variables. The completeness of the families of PDFs is equivalent to the invertibility of operators using these PDFs as kernel functions. Invertibility permits the nontrivial transformation of semi-parametric censored dynamic panel data models into a valid semi-parametric family of PDFs of observables. Then, identification can be achieved under the MLE framework. The dynamic tobit models and dynamic lognormal hurdle models with two common types of DGPs satisfy these completeness conditions. This identification leads to the proposed sieve MLE, which is consistent and asymptotically normal. The advantage of the proposed approach is that it does not rely on the availability of initial period data, provides average partial effects, and requires only two-period data. In addition, this semi-parametric method allows for time dummies, nonlinear functions of state dependence Y_{it-1}

such as quadratics or interaction terms, and parametric heteroskedasticity. These features make the sieve MLE desirable in semi-parametric censored dynamic panel data models for microeconomic applications.

Appendix

A. Proof of Lemma 2.1

Proof: First, we have shown $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0} = f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}}$. Next, given (x_{it}, y_{it-1}) , define integral operators

$$L_{f_{Y_{it}^+ | X_{it}, Y_{it-1}, X_{it-1}}} : \mathcal{L}^2(\tilde{\mathcal{Y}}_{it}) \rightarrow \mathcal{L}^2(\mathcal{X}_{it-1}) \text{ with} \quad (37)$$

$$(L_{f_{Y_{it}^+ | X_{it}, Y_{it-1}, X_{it-1}}} h)(x_{it-1}) = \int f_{Y_{it}^+ | X_{it}, Y_{it-1}, X_{it-1}}(y_{it}^+, x_{it}, y_{it-1}, x_{it-1}) h(y_{it}^+) dy_{it}^+,$$

$$\tilde{L}_{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0}}} : \mathcal{L}^2(\tilde{\mathcal{Y}}_{it}) \rightarrow \mathcal{L}^2(\mathcal{U}_{it}, \omega) \text{ with} \quad (38)$$

$$(\tilde{L}_{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0}}} h)(u_{it}) = \int f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0}}(y_{it}^+ | x_{it}, y_{it-1}, u_{it}) h(y_{it}^+) dy_{it}^+,$$

$$L_{f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}}} : \mathcal{L}^2(\mathcal{U}_{it}, \omega) \rightarrow \mathcal{L}^2(\mathcal{X}_{it-1}) \text{ with} \quad (39)$$

$$(L_{f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}}} h)(x_{it-1}) = \int \frac{f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}}{\omega(u_{it})} h(u_{it}) \omega(u_{it}) du_{it}.$$

For each $h \in \mathcal{L}^2(\tilde{\mathcal{Y}}_{it})$.

$$\begin{aligned} & \left(L_{f_{Y_{it}^+ | X_{it}, Y_{it-1}, X_{it-1}}} \right) (h) (x_{it-1}) \\ &= \int_{\tilde{\mathcal{Y}}_{it}} f_{Y_{it}^+ | X_{it}, Y_{it-1}, X_{it-1}} h(y_{it}^+) dy_{it}^+ \\ &= \int_{\tilde{\mathcal{Y}}_{it}} \left(\int_{\mathcal{U}_{it}} f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0}} \frac{f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}}{\omega(u_{it})} \omega(u_{it}) du_{it} \right) h(y_{it}^+) dy_{it}^+ \\ &= \int_{\mathcal{U}_{it}} \frac{f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}}{\omega(u_{it})} \left(\int_{\tilde{\mathcal{Y}}_{it}} f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0}} h(y_{it}^+) dy_{it}^+ \right) \omega(u_{it}) du_{it} \\ &= \left(L_{f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}}} \tilde{L}_{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0}}} \right) (h) (x_{it-1}), \end{aligned}$$

based on Eq. (12). Because this derivation holds for arbitrary h , this amounts to the operator relationship

$$\underbrace{L_{f_{Y_{it}^+, X_{it}, Y_{it-1}, X_{it-1}}}}_{\text{Assumption 2.4(i)}} = L_{f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}} \underbrace{\tilde{L}_{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0}}}_{\text{Assumption 2.4(ii)}}.$$

Combining the condition $f_{X_{it}, Y_{it-1}, X_{it-1}} > 0$ and Assumption 2.4(i) results in $\{f_{Y_{it}^+, X_{it}, Y_{it-1}, X_{it-1}} : x_{it-1} \in \mathcal{X}_{it-1}\}$, is complete over $\mathcal{L}^2(\tilde{\mathcal{Y}}_{it})$ and then $L_{f_{Y_{it}^+, X_{it}, Y_{it-1}, X_{it-1}}}$ is invertible. In addition, because Assumption 2.4(ii) ensures the operator $\tilde{L}_{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0}}$ invertible, the operator relationship implies that the invertibility of the operator $L_{f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}}$, i.e., $\{\frac{f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}}{\omega(u_{it})} : x_{it-1} \in \mathcal{X}_{it-1}\}$ is complete over $\mathcal{L}^2(\mathcal{U}_{it}, \omega)$.

Suppose that the parameter θ_0 is not locally identifiable. Then, there exists $\theta_k \neq \theta_0$ and $\theta_k \mapsto \theta_0$ such that $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0} = f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_k}$. Using the definition of $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}$ and $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_k}$,

$$f_{Y_{it}^+, X_{it}, Y_{it-1}, X_{it-1}} = \int f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0} du_{it}, \quad (40)$$

$$f_{Y_{it}^+, X_{it}, Y_{it-1}, X_{it-1}} = \int f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_k} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_k} du_{it}. \quad (41)$$

By subtracting Eq. (41) from Eq. (40), it follows that

$$\begin{aligned} 0 &= \int f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_k} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_k} - f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0} du_{it}, \\ &= \int f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_k} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_k} - f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_k} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0} du_{it} \\ &\quad + \int f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_k} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0} - f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0} du_{it}, \\ &= \int \underbrace{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_k}}_{\text{Assumption 2.3}} (f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_k} - f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}) du_{it} \\ &\quad + \int (f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_k} - f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0}) \underbrace{f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}}_{\text{Assumption 2.4(i) \& (ii)}} du_{it}. \end{aligned} \quad (42)$$

Plugging the relation $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0} = f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_k}$ into the above equation yields

$$0 = \int \left(f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_k} - f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0} \right) \frac{f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}}{\omega(u_{it})} \omega(u_{it}) du_{it},$$

for all x_{it-1} in \mathcal{X}_{it-1} . Because Assumptions 2.4(i) & (ii) implies that $\left\{ \frac{f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}}{\omega(u_{it})} : x_{it-1} \in \mathcal{X}_{it-1} \right\}$ is complete over $\mathcal{L}^2(\mathcal{U}_{it}, \omega)$, we obtain $f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_k} = f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0}$ for $\theta_k \neq \theta_0$ and $\theta_k \mapsto \theta_0$. This contradicts to the local identifiability of θ_0 in $f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta}$, proving the lemma. Q.E.D.

B. Proof of Lemma 2.2

Proof: Because $f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta}$ is correctly specified at θ_0 by Lemma 2.1, $\tilde{f}_{X_{it}, Y_{it-1}, X_{it-1}; \theta}$ is also correctly specified at θ_0 after integrating out. On the other hand, denote two integral kernels as $K_{A; \theta_0}(x_{it}, y_{it-1}, x_{it-1}, u_{it}) \equiv \frac{1}{\omega(u_{it})} \frac{\partial}{\partial \theta} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}$ and $K_{B; \theta_0}(y_{it}^+, x_{it}, y_{it-1}, u_{it}) \equiv \frac{\partial}{\partial \theta} f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0}$. Divide Eq. (42) by $\theta - \theta_0 \neq 0$ and rewrite it as follows:

$$\begin{aligned} 0 = & \int f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta} \frac{1}{\omega(u_{it})} \frac{f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta} - f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}}{\theta - \theta_0} \omega(u_{it}) du_{it} \\ & + \int \frac{f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta} - f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0}}{\theta - \theta_0} \frac{f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}}{\omega(u_{it})} \omega(u_{it}) du_{it}. \end{aligned}$$

If $\theta \mapsto \theta_0$ then the above equation implies

$$\begin{aligned} 0 = & \int f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0} K_{A; \theta_0}(x_{it}, y_{it-1}, x_{it-1}, u_{it}) \omega(u_{it}) du_{it} \\ & + \int K_{B; \theta_0}(y_{it}^+, x_{it}, y_{it-1}, u_{it}) \frac{f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}}{\omega(u_{it})} \omega(u_{it}) du_{it}. \end{aligned} \quad (43)$$

This equation can be used to establish an operator relationship. For each given (x_{it}, y_{it-1}) , define integral operators as follows

$$L_{K_{A; \theta_0}} : \mathcal{L}^2(\mathcal{U}_{it}, \omega) \rightarrow \mathcal{L}^2(\mathcal{X}_{it-1}) \text{ with} \quad (44)$$

$$(L_{K_{A; \theta_0}} h)(x_{it-1}) = \int \frac{1}{\omega(u_{it})} \frac{\partial}{\partial \theta} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}(x_{it}, y_{it-1}, x_{it-1}, u_{it}) h(u_{it}) \omega(u_{it}) du_{it},$$

$$L_{K_{B; \theta_0}} : \mathcal{L}^2(\tilde{\mathcal{Y}}_{it}) \rightarrow \mathcal{L}^2(\mathcal{U}_{it}, \omega) \text{ with} \quad (45)$$

$$(L_{K_{B; \theta_0}} h)(u_{it}) = \int \frac{\partial}{\partial \theta} f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta_0}(y_{it}^+ | x_{it}, y_{it-1}, u_{it}) h(y_{it}^+) dy_{it}^+.$$

Set $h \in \mathcal{L}^2(\mathcal{X}_{it-1})$. Given each (x_{it}, y_{it-1}) ,

$$\begin{aligned}
& \left(L_{K_{A;\theta_0}} \tilde{L}_{f_{Y_{it}^+|X_{it},Y_{it-1},U_{it};\theta_0}} \right) (h) (x_{it-1}) \\
&= \int_{\mathcal{U}_{it}} K_{A;\theta_0}(x_{it}, y_{it-1}, x_{it-1}, u_{it}) \left(\int_{\mathcal{Y}_{it}} f_{Y_{it}^+|X_{it},Y_{it-1},U_{it};\theta_0} h(y_{it}^+) dy_{it}^+ \right) \omega(u_{it}) du_{it} \\
&= \int_{\mathcal{Y}_{it}} \left(\int_{\mathcal{U}_{it}} f_{Y_{it}^+|X_{it},Y_{it-1},U_{it};\theta_0} K_{A;\theta_0}(x_{it}, y_{it-1}, x_{it-1}, u_{it}) \omega(u_{it}) du_{it} \right) h(y_{it}^+) dy_{it}^+ \\
&= - \int_{\mathcal{Y}_{it}} \left(\int_{\mathcal{U}_{it}} K_{B;\theta_0}(y_{it}^+, x_{it}, y_{it-1}, u_{it}) \frac{f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta_0}}{\omega(u_{it})} \omega(u_{it}) du_{it} \right) h(y_{it}^+) dy_{it}^+ \\
&= - \int_{\mathcal{U}_{it}} \frac{f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta_0}}{\omega(u_{it})} \left(\int_{\mathcal{Y}_{it}} K_{B;\theta_0}(y_{it}^+, x_{it}, y_{it-1}, u_{it}) h(y_{it}^+) dy_{it}^+ \right) \omega(u_{it}) du_{it} \\
&= - \left(L_{f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta_0}} L_{K_{B;\theta_0}} \right) (h) (x_{it-1})
\end{aligned}$$

where we have used (i) an interchange of the order of integration (justified by Fubini's theorem), (ii) Eq. (43), (iii) the definitions of these operators in Eqs. (38), (39), (44), and (45). This derivation yields the following operator relationship

$$L_{K_{A;\theta_0}} \underbrace{\tilde{L}_{f_{Y_{it}^+|X_{it},Y_{it-1},U_{it};\theta_0}}}_{\text{Assumptions 2.4(ii)}} + \underbrace{L_{f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta_0}}}_{\text{Lemma 2.1}} \underbrace{L_{K_{B;\theta_0}}}_{\text{Assumptions 2.5}} = 0. \quad (46)$$

Whereas Assumptions 2.4(i) & (ii) imply that $\tilde{L}_{f_{Y_{it}^+|X_{it},Y_{it-1},U_{it};\theta_0}}$ and $L_{f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta_0}}$ are invertible,¹⁵ Assumption 2.5 guarantees that $L_{K_{B;\theta_0}}$ is invertible. Because the operators other than $L_{K_{A;\theta_0}}$ in Eq. (46) are all invertible, the integral operator $L_{K_{A;\theta_0}}$ is also invertible. This implies that the family of its corresponding kernel functions $\{ \frac{1}{\omega(u_{it})} \frac{\partial}{\partial \theta} f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta_0} : x_{it-1} \in \mathcal{X}_{it-1} \}$ is complete over $\mathcal{L}^2(\mathcal{U}_{it}, \omega)$.

Suppose θ_0 is not locally identifiable in $\tilde{f}_{X_{it},Y_{it-1},X_{it-1};\theta}$. This implies that there exists $\theta_k \neq \theta_0$ and $\theta_k \mapsto \theta_0$ such that $\tilde{f}_{X_{it},Y_{it-1},X_{it-1};\theta_k}(x_{it}, y_{it-1}, x_{it-1}) = \tilde{f}_{X_{it},Y_{it-1},X_{it-1};\theta_0}(x_{it}, y_{it-1}, x_{it-1})$. This implies that $\int f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta_k} du_{it} = \int f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta_0} du_{it}$ for each θ_k . It follows that for each θ_k

$$\int \frac{1}{\omega(u_{it})} \left(\frac{f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta_k} - f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta_0}}{\theta_k - \theta_0} \right) \omega(u_{it}) du_{it} = 0 \quad \text{for all } x_{it-1}.$$

¹⁵ Assumptions 2.4(i) & (ii) imply that the invertibility of $L_{f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta_0}}$ is in the proof of Lemma 2.1.

If $\theta_k \mapsto \theta_0$, the equation becomes

$$\int \left(\frac{1}{\omega(u_{it})} \frac{\partial}{\partial \theta} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0} \right) \omega(u_{it}) du_{it} = 0 \quad \text{for all } x_{it-1}. \quad (47)$$

Because $\mathcal{L}^2(\mathcal{U}_{it}, \omega)$ contains the constant function, Eq. (47) is in contradiction with the completeness of $\left\{ \frac{1}{\omega(u_{it})} \frac{\partial}{\partial \theta} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0} : x_{it-1} \in \mathcal{X}_{it-1} \right\}$ over $\mathcal{L}^2(\mathcal{U}_{it}, \omega)$. Therefore, under Assumptions 2.3-2.5, θ_0 is locally identifiable. *Q.E.D.*

C. Proof of Lemma 2.3

Before proving Lemma 2.3, consider the following result as the cornerstone of the proof of Lemma 2.3.

Lemma C.1. *Under Assumptions 2.3-2.6, the family of functions $\left\{ \frac{1}{\omega(u_{it})} \frac{\partial}{\partial x_{it-1}} f_{U_{it}|X_{it}, Y_{it-1}, X_{it-1}} : x_{it-1} \in \mathcal{X}_{it-1} \right\}$ is complete over $\mathcal{L}^2(\mathcal{U}_{it}, \omega)$.*

Proof: In a similar manner to Eq. (12), write the conditional version of Eq. (12) for $\theta = \theta_0$,

$$f_{Y_{it}^+|X_{it}, Y_{it-1}, X_{it-1}} = \int f_{Y_{it}^+|X_{it}, Y_{it-1}, U_{it}} f_{U_{it}|X_{it}, Y_{it-1}, X_{it-1}} du_{it}. \quad (48)$$

Taking derivative with respect to X_{it-1} results in

$$\frac{\partial}{\partial x_{it-1}} f_{Y_{it}^+|X_{it}, Y_{it-1}, X_{it-1}} = \int f_{Y_{it}^+|X_{it}, Y_{it-1}, U_{it}} \frac{\partial}{\partial x_{it-1}} f_{U_{it}|X_{it}, Y_{it-1}, X_{it-1}} du_{it}. \quad (49)$$

Set $\kappa_1 = \frac{\partial}{\partial x_{it-1}} f_{Y_{it}^+|X_{it}, Y_{it-1}, X_{it-1}}$ and $\phi = \frac{\partial}{\partial x_{it-1}} f_{U_{it}|X_{it}, Y_{it-1}, X_{it-1}}$. For each (x_{it}, y_{it-1}) , define operators

$$L_{\kappa_1} : \mathcal{L}^2(\tilde{\mathcal{Y}}_{it}) \rightarrow \mathcal{L}^2(\mathcal{X}_{it-1}) \text{ with } (L_{\kappa_1} h)(x_{it-1}) = \int \kappa_1(y_{it}^+, x_{it}, y_{it-1}, x_{it-1}) h(y_{it}^+) dy_{it}^+,$$

$$L_{\phi} : \mathcal{L}^p(\mathcal{U}_{it}, \omega) \rightarrow \mathcal{L}^2(\mathcal{X}_{it-1}) \text{ with}$$

$$(L_{\phi} h)(x_{it-1}) = \int \frac{1}{\omega(u_{it})} \phi(u_{it}, x_{it}, y_{it-1}, x_{it-1}) h(u_{it}) \omega(u_{it}) du_{it}.$$

For $h \in \mathcal{L}^2(\tilde{\mathcal{Y}}_{it})$.

$$\begin{aligned}
& (L_{\kappa_1})(h)(x_{it-1}) \\
&= \int_{\tilde{\mathcal{Y}}_{it}} \kappa_1(y_{it}^+, x_{it}, y_{it-1}, x_{it-1}) h(y_{it}^+) dy_{it}^+ \\
&= \int_{\tilde{\mathcal{Y}}_{it}} \left(\int_{\mathcal{U}_{it}} f_{Y_{it}^+|X_{it}, Y_{it-1}, U_{it}} \phi(u_{it}, x_{it}, y_{it-1}, x_{it-1}) du_{it} \right) h(y_{it}^+) y_{it}^+ \\
&= \int_{\mathcal{U}_{it}} \frac{1}{\omega(u_{it})} \phi(u_{it}, x_{it}, y_{it-1}, x_{it-1}) \left(\int_{\mathcal{Y}_{it}} f_{Y_{it}^+|X_{it}, Y_{it-1}, U_{it}} h(y_{it}) dy_{it} \right) \omega(u_{it}) du_{it} \\
&= \left(L_\phi \tilde{L}_{f_{Y_{it}^+|X_{it}, Y_{it-1}, U_{it}; \theta_0}} \right) (h)(x_{it-1}),
\end{aligned}$$

where Eq. (38) defines the operator $\tilde{L}_{f_{Y_{it}^+|X_{it}, Y_{it-1}, U_{it}; \theta_0}}$. With the definitions of the operators, this equation can be rewritten as an operator relationship

$$L_{\kappa_1} = L_\phi \tilde{L}_{f_{Y_{it}^+|X_{it}, Y_{it-1}, U_{it}; \theta_0}}. \quad (50)$$

Assumptions 2.4(ii) and 2.6 guarantee the invertibility of the operators $\tilde{L}_{f_{Y_{it}^+|X_{it}, Y_{it-1}, U_{it}; \theta_0}}$ and L_{κ_1} , respectively. Applying this invertibility to Eq. (50) results in the invertibility of L_ϕ . Thus, the family $\{\frac{1}{\omega(u_{it})} \phi(u_{it}, x_{it}, y_{it-1}, x_{it-1}) : x_{it-1} \in \mathcal{X}_{it-1}\}$ is complete over $\mathcal{L}^2(\mathcal{U}_{it}, \omega)$ for each x_{it}, y_{it-1} . *Q.E.D.*

Proof of Lemma 2.3: First, $f_{X_{it}, Y_{it-1}, X_{it-1}; \theta}$ is correctly specified at θ_0 because by Lemma 2.2, $\tilde{f}_{X_{it}, Y_{it-1}, X_{it-1}; \theta}$ is correctly specified at θ_0 . Suppose that θ_0 is not locally identifiable in the observable joint density function $f_{X_{it}, Y_{it-1}, X_{it-1}; \theta}$. There exists $\theta_k \neq \theta_0$ and $\theta_k \mapsto \theta_0$ such that, $f_{X_{it}, Y_{it-1}, X_{it-1}; \theta_k} = f_{X_{it}, Y_{it-1}, X_{it-1}; \theta_0}$. This implies that

$$\frac{\tilde{f}_{X_{it}, Y_{it-1}, X_{it-1}; \theta_k}}{\int \int \int \tilde{f}_{X_{it}, Y_{it-1}, X_{it-1}; \theta_k} dx_{it} dy_{it-1} dx_{it-1}} = \frac{\tilde{f}_{X_{it}, Y_{it-1}, X_{it-1}; \theta_0}}{1} = f_{X_{it}, Y_{it-1}, X_{it-1}}. \quad (51)$$

This equation can be expressed as

$$\frac{\int f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_k} du_{it}}{f_{X_{it}, Y_{it-1}, X_{it-1}}} = \int \int \int \tilde{f}_{X_{it}, Y_{it-1}, X_{it-1}; \theta_k} dx_{it} dy_{it-1} dx_{it-1}. \quad (52)$$

The multiple integral in the RHS of Eq. (52) only depends on the parameter θ_k , and is

independent of x_{it-1} . This suggests that given $x_{1t-1} \neq x_{2t-1}$,

$$\int \frac{f_{X_{it}, Y_{it-1}, X_{1t-1}, U_{it}; \theta_k}}{f_{X_{it}, Y_{it-1}, X_{1t-1}}} du_{it} = \int \frac{f_{X_{it}, Y_{it-1}, X_{2t-1}, U_{it}; \theta_k}}{f_{X_{it}, Y_{it-1}, X_{2t-1}}} du_{it}.$$

If $\theta_k \mapsto \theta_0$, this yields

$$0 = \int (f_{U_{it}|X_{it}, Y_{it-1}, X_{1t-1}} - f_{U_{it}|X_{it}, Y_{it-1}, X_{2t-1}}) du_{it}$$

Divide the equation by $X_{1t-1} - X_{2t-1}$ and let $X_{1t-1} - X_{2t-1} \mapsto 0$. This equation then changes into

$$\begin{aligned} 0 &= \int \frac{\partial}{\partial x_{it-1}} f_{U_{it}|X_{it}, Y_{it-1}, X_{1t-1}} du_{it} \\ &= \int \frac{1}{\omega(u_{it})} \frac{\partial}{\partial x_{it-1}} f_{U_{it}|X_{it}, Y_{it-1}, X_{1t-1}} \omega(u_{it}) du_{it}, \end{aligned}$$

which contradicts the completeness in Lemma C.1. Therefore, the parameter θ_0 is locally identifiable in the observable joint density function $f_{X_{it}, Y_{it-1}, X_{it-1}; \theta}$. *Q.E.D.*

D. Identification in the Discrete Case

This section presents a simple case in which the observed variables $Y_{it}, X_{it}, Y_{it-1}, X_{it-1}$ and the unobserved covariate U_{it} are all discrete. This section shows how to use the identification techniques in Theorem 2.1 for this discrete case. For simplicity, assume that the variables Y_{it}^+, X_{it-1} and U_{it} have the same size J (i.e., $Y_{it}^+, X_{it-1}, U_{it} \in \{1, 2, \dots, J\}$). For this setting, the integral operators used previously can be represented by J -by- J matrices. The idea of using the identification strategy in the discrete case for ease of exposition is because a complete integral operators is associated with an invertible matrix.¹⁶

¹⁶If $y, u \in \{1, 2\}$ and $\int_{\mathcal{U}} h(u) f(y|u) du = 0$, then the condition is equivalent to

$$\begin{bmatrix} f_{y|u}(1|1) & f_{y|u}(1|2) \\ f_{y|u}(2|1) & f_{y|u}(2|2) \end{bmatrix} \begin{bmatrix} h(1) \\ h(2) \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

The function h can be uniquely determined as $h = 0$ iff the first matrix representing $f_{y|u}$ is invertible.

Equation (12) in the discrete case is

$$f_{Y_{it}^+, X_{it}, Y_{it-1}, X_{it-1}} = \sum_{U_{it}=1}^J f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta}. \quad (53)$$

Given (x_{it}, y_{it-1}) , define J -by- J matrices

$$\begin{aligned} M_{f_{Y_{it}^+, x_{it}, y_{it-1}, X_{it-1}}} &= \left[f_{Y_{it}^+, X_{it}, Y_{it-1}, X_{it-1}}(Y_{it}^+, x_{it}, y_{it-1}, X_{it-1}) \right]_{y_{it}^+, x_{it-1}} \\ L_{f_{Y_{it}^+ | x_{it}, y_{it-1}, U_{it}; \theta}} &= \left[f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta}(Y_{it}^+, x_{it}, y_{it-1}, U_{it}) \right]_{y_{it}^+, u_{it}} \\ M_{f_{x_{it}, y_{it-1}, X_{it-1}, U_{it}; \theta}} &= \left[f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta}(x_{it}, y_{it-1}, X_{it-1}, U_{it}) \right]_{u_{it}, x_{it-1}}. \end{aligned}$$

Rewrite the equality (53) in terms of these matrices as follows:

$$\underbrace{M_{f_{Y_{it}^+, x_{it}, y_{it-1}, X_{it-1}}}}_{\text{Observed from Data}} = \underbrace{L_{f_{Y_{it}^+ | x_{it}, y_{it-1}, U_{it}; \theta}}}_{\text{Model Specification}} M_{f_{x_{it}, y_{it-1}, X_{it-1}, U_{it}; \theta}}. \quad (54)$$

Assumption 2.3 implies that the square matrix $L_{f_{Y_{it}^+ | x_{it}, y_{it-1}, U_{it}; \theta}}$ is invertible, leading to

$$M_{f_{x_{it}, y_{it-1}, X_{it-1}, U_{it}; \theta}} = \left(L_{f_{Y_{it}^+ | x_{it}, y_{it-1}, U_{it}; \theta}} \right)^{-1} M_{f_{Y_{it}^+, x_{it}, y_{it-1}, X_{it-1}}}. \quad (55)$$

As discussed earlier, it is necessary to ensure that $M_{f_{x_{it}, y_{it-1}, X_{it-1}, U_{it}; \theta}}$ is identifiable at θ_0 . According to the proof of Lemma 2.1, there are two steps for identifiability. First, given (x_{it}, y_{it-1}) , define

$$\begin{aligned} L_{f_{Y_{it}^+, x_{it}, y_{it-1}, X_{it-1}}} &= \left[f_{Y_{it}^+, X_{it}, Y_{it-1}, X_{it-1}}(Y_{it}^+, x_{it}, y_{it-1}, X_{it-1}) \right]_{x_{it-1}, y_{it}^+} \\ \tilde{L}_{f_{Y_{it}^+ | x_{it}, y_{it-1}, U_{it}; \theta}} &= \left[f_{Y_{it}^+ | X_{it}, Y_{it-1}, U_{it}; \theta}(Y_{it}^+, x_{it}, y_{it-1}, U_{it}) \right]_{u_{it}, y_{it}^+} \\ L_{f_{x_{it}, y_{it-1}, X_{it-1}, U_{it}; \theta}} &= \left[f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta}(x_{it}, y_{it-1}, x_{it-1}, U_{it}) \right]_{x_{it-1}, u_{it}}. \end{aligned}$$

Equality (53) can then be expressed by these matrices as follows:

$$\underbrace{L_{f_{Y_{it}^+, x_{it}, y_{it-1}, X_{it-1}}}}_{\text{Assumption 2.4(i)}} = L_{f_{x_{it}, y_{it-1}, X_{it-1}, U_{it}; \theta_0}} \underbrace{\tilde{L}_{f_{Y_{it}^+ | x_{it}, y_{it-1}, U_{it}; \theta_0}}}_{\text{Assumption 2.4(ii)}}. \quad (56)$$

Notice that in this simple case, $L_{f_{Y_{it}^+|x_{it},y_{it-1},X_{it-1}}} = M_{f_{Y_{it}^+|x_{it},y_{it-1},X_{it-1}}}^T$, $\tilde{L}_{f_{Y_{it}^+|x_{it},y_{it-1},U_{it};\theta}} = L_{f_{Y_{it}^+|x_{it},y_{it-1},U_{it};\theta}}^T$, and $L_{f_{x_{it},y_{it-1},X_{it-1},U_{it};\theta}} = M_{f_{x_{it},y_{it-1},X_{it-1},U_{it};\theta}}^T$ which we may not have for a general continuous case. The matrix notations used here are based on integral operators in the proofs of lemmas. Assumption 2.4 makes $L_{f_{x_{it},y_{it-1},X_{it-1},U_{it};\theta_0}}$ invertible. Hence, its transpose $M_{f_{x_{it},y_{it-1},X_{it-1},U_{it};\theta_0}}$ is also invertible. Then, suppose that there exists $\theta_k \neq \theta_0$ and $\theta_k \mapsto \theta_0$ such that

$$M_{f_{x_{it},y_{it-1},X_{it-1},U_{it};\theta_k}} = M_{f_{x_{it},y_{it-1},X_{it-1},U_{it};\theta_0}}. \quad (57)$$

Following the derivation in Eq. (42), we have a matrix expression

$$\begin{aligned} 0 &= L_{f_{Y_{it}^+|x_{it},y_{it-1},U_{it};\theta_k}} \left(M_{f_{x_{it},y_{it-1},X_{it-1},U_{it};\theta_k}} - M_{f_{x_{it},y_{it-1},X_{it-1},U_{it};\theta_0}} \right) \\ &\quad + \left(L_{f_{Y_{it}^+|x_{it},y_{it-1},U_{it};\theta_k}} - L_{f_{Y_{it}^+|x_{it},y_{it-1},U_{it};\theta_0}} \right) M_{f_{x_{it},y_{it-1},X_{it-1},U_{it};\theta_0}} \end{aligned}$$

The invertibility of $M_{f_{x_{it},y_{it-1},X_{it-1},U_{it};\theta_0}}$ and Eq. (57) implies that $L_{f_{Y_{it}^+|x_{it},y_{it-1},U_{it};\theta}}$ is not identifiable at θ_0 , which is a contradiction.

Set $J \times 1$ -vector $\mathbf{J}_1 = (1, 1, \dots, 1)^T$. Integrating out the unobserved covariate in the discrete case leads to $\mathbf{J}_1^T M_{f_{x_{it},y_{it-1},X_{it-1},U_{it};\theta}}$. Suppose that there exists $\theta_k \neq \theta_0$ and $\theta_k \mapsto \theta_0$ such that $\mathbf{J}_1^T M_{f_{x_{it},y_{it-1},X_{it-1},U_{it};\theta_k}} = \mathbf{J}_1^T M_{f_{x_{it},y_{it-1},X_{it-1},U_{it};\theta_0}}$. It then follows that

$$0 = \mathbf{J}_1^T \left(\frac{M_{f_{x_{it},y_{it-1},X_{it-1},U_{it};\theta_k}} - M_{f_{x_{it},y_{it-1},X_{it-1},U_{it};\theta_0}}}{\theta_k - \theta_0} \right) \quad (58)$$

If $\theta \mapsto \theta_0$, the above equation implies $0 = \mathbf{J}_1^T M_{\frac{\partial}{\partial \theta} f_{x_{it},y_{it-1},X_{it-1},U_{it};\theta_0}}$, where

$$M_{\frac{\partial}{\partial \theta} f_{x_{it},y_{it-1},X_{it-1},U_{it};\theta}} = \left[\frac{\partial}{\partial \theta} f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta}(x_{it}, y_{it-1}, X_{it-1}, U_{it}) \right]_{u_{it}, x_{it-1}}.$$

Rewrite Eq. (43) in the discrete case as

$$\begin{aligned} 0 &= \sum_{U_{it}=1}^J f_{Y_{it}^+|X_{it},Y_{it-1},U_{it};\theta_0} \frac{\partial}{\partial \theta} f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta_0} \\ &\quad + \sum_{U_{it}=1}^J \frac{\partial}{\partial \theta} f_{Y_{it}^+|X_{it},Y_{it-1},U_{it};\theta_0} f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta_0}. \end{aligned} \quad (59)$$

This leads to the following matrix expression

$$M_{\frac{\partial}{\partial \theta} f_{x_{it}, y_{it-1}, X_{it-1}, U_{it}; \theta_0}} \underbrace{\tilde{L}_{f_{Y_{it}^+ | x_{it}, y_{it-1}, U_{it}; \theta_0}}}_{\text{Assumptions 2.4(ii)}} + \underbrace{L_{f_{x_{it}, y_{it-1}, X_{it-1}, U_{it}; \theta_0}}}_{\text{Lemma 2.1}} \underbrace{M_{\frac{\partial}{\partial \theta} f_{Y_{it}^+ | x_{it}, y_{it-1}, U_{it}; \theta_0}}}_{\text{Assumptions 2.5}} = 0, \quad (60)$$

where $M_{\frac{\partial}{\partial \theta} f_{Y_{it}^+ | x_{it}, y_{it-1}, U_{it}; \theta_0}} = \left[\frac{\partial}{\partial \theta} f_{Y_{it}^+ | x_{it}, y_{it-1}, U_{it}; \theta_0} \right]_{u_{it}, y_{it}^+}$. Applying assumptions to Eq. (60) shows that $M_{\frac{\partial}{\partial \theta} f_{x_{it}, y_{it-1}, X_{it-1}, U_{it}; \theta_0}}$ is invertible, which contradicts $0 = \mathbf{J}_1^T M_{\frac{\partial}{\partial \theta} f_{x_{it}, y_{it-1}, X_{it-1}, U_{it}; \theta_0}}$.

Finally, the normalization in the discrete case is equivalent to

$$\mathbf{V}_{f_{x_{it}, y_{it-1}, X_{it-1}, \theta}} \equiv \frac{\mathbf{J}_1^T M_{f_{x_{it}, y_{it-1}, X_{it-1}, U_{it}; \theta}}}{\sum_{x_{it}=1}^{J_{x_{it}}} \sum_{y_{it-1}=1}^{J_{y_{it-1}}} \left(\mathbf{J}_1^T M_{f_{x_{it}, y_{it-1}, X_{it-1}, U_{it}; \theta}} \mathbf{J} \right)}, \quad (61)$$

where $J_{x_{it}}$ and $J_{y_{it-1}}$ represent the sizes of the discrete variables x_{it} and y_{it-1} , respectively. Suppose the normalization step does not lead to local identifiability at θ_0 . This implies that there exists $\theta_k \neq \theta_0$ and $\theta_k \mapsto \theta_0$ such that

$$\frac{\mathbf{J}_1^T M_{f_{x_{it}, y_{it-1}, X_{it-1}, U_{it}; \theta_k}}}{\sum_{x_{it}=1}^{J_{x_{it}}} \sum_{y_{it-1}=1}^{J_{y_{it-1}}} \left(\mathbf{J}_1^T M_{f_{x_{it}, y_{it-1}, X_{it-1}, U_{it}; \theta_k}} \mathbf{J} \right)} = \frac{\mathbf{J}_1^T M_{f_{x_{it}, y_{it-1}, X_{it-1}, U_{it}; \theta_0}}}{1} = \mathbf{V}_{f_{x_{it}, y_{it-1}, X_{it-1}, \theta_0}} \quad (62)$$

Rearrange the term

$$\left(\mathbf{J}_1^T M_{f_{x_{it}, y_{it-1}, X_{it-1}, U_{it}; \theta_k}} \right) ./ \mathbf{V}_{f_{x_{it}, y_{it-1}, X_{it-1}, \theta_0}} = \left(\sum_{x_{it}=1}^{J_{x_{it}}} \sum_{y_{it-1}=1}^{J_{y_{it-1}}} \left(\mathbf{J}_1^T M_{f_{x_{it}, y_{it-1}, X_{it-1}, U_{it}; \theta_k}} \mathbf{J} \right) \right) \mathbf{J}_1,$$

where the notation $./$ divides two $1 \times J$ -vectors element-wise. The right-hand side of this equation is constant in x_{it-1} . Hence, if $x_{1t-1} \neq x_{2t-1}$, we have

$$\left(\mathbf{J}_1^T M_{f_{x_{it}, y_{it-1}, X_{1t-1}, U_{it}; \theta_k}} \right) ./ \mathbf{V}_{f_{x_{it}, y_{it-1}, X_{1t-1}, \theta_0}} = \left(\mathbf{J}_1^T M_{f_{x_{it}, y_{it-1}, X_{2t-1}, U_{it}; \theta_k}} \right) ./ \mathbf{V}_{f_{x_{it}, y_{it-1}, X_{2t-1}, \theta_0}}.$$

Using Eq. (55), rewrite this equation as

$$0 = \mathbf{J}_1^T \left(L_{f_{Y_{it}^+ | x_{it}, y_{it-1}, U_{it}; \theta_k}} \right)^{-1} \left(M_{f_{Y_{it}^+ | x_{it}, y_{it-1}, X_{1t-1}}} - M_{f_{Y_{it}^+ | x_{it}, y_{it-1}, X_{2t-1}}} \right) ./ \mathbf{V}_{f_{x_{it}, y_{it-1}, X_{2t-1}, \theta_0}}.$$

Denote $M_{f_{Y_{it}^+ | x_{it}, y_{it-1}, \Delta X_{it-1}}}$ as a matrix of the difference of $f_{Y_{it}^+ | x_{it}, y_{it-1}, X_{it-1}}$ with respect to

X_{it-1} . If $\theta_k \mapsto \theta_0$, then

$$\begin{aligned} 0 &= \mathbf{J}_1^T \left(L_{f_{Y_{it}^+|x_{it},y_{it-1},U_{it};\theta_0}} \right)^{-1} \left(M_{f_{Y_{it}^+|x_{it},y_{it-1},X_{1t-1}}} - M_{f_{Y_{it}^+|x_{it},y_{it-1},X_{2t-1}}} \right) \\ &\equiv \underbrace{\mathbf{J}_1^T \left(L_{f_{Y_{it}^+|x_{it},y_{it-1},U_{it};\theta_0}} \right)^{-1}}_{\text{Assumption 2.3}} \underbrace{M_{f_{Y_{it}^+|x_{it},y_{it-1},\Delta X_{it-1}}}}_{\text{Assumption 2.6}}, \end{aligned}$$

where $M_{f_{Y_{it}^+|x_{it},y_{it-1},X_{it-1}}} \equiv \left[f_{Y_{it}^+|X_{it},Y_{it-1},X_{it-1}}(Y_{it}^+|x_{it},y_{it-1},X_{it-1}) \right]_{y_{it}^+,x_{it-1}}$. This contradicts the invertibility under Assumptions 2.3 and 2.6, showing that the density function is still locally identifiable at θ_0 .

E. Proof of a Sufficient Condition of the Uniqueness of $K(\theta)$

Proof: Start with a scalar θ , and combine Eqs. (16) and (17),

$$f_{X_{it},Y_{it-1},X_{it-1};\theta} \equiv \frac{\int f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta} du_{it}}{\int \int \int \int f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta} du_{it} dx_{it} dy_{it-1} dx_{it-1}}. \quad (63)$$

and recall Eq. (43),

$$\begin{aligned} 0 &= \int f_{Y_{it}^+|X_{it},Y_{it-1},U_{it};\theta_0} \frac{\partial}{\partial \theta} f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta} \Big|_{\theta=\theta_0} du_{it} \\ &\quad + \int \frac{\partial}{\partial \theta} f_{Y_{it}^+|X_{it},Y_{it-1},U_{it};\theta_0} f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta_0} du_{it}. \end{aligned} \quad (64)$$

It follows that $\frac{\partial}{\partial \theta} f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta} \Big|_{\theta=\theta_0}$ is implicitly defined in Eq. (64) by

$$\frac{\partial}{\partial \theta} f_{X_{it},Y_{it-1},X_{it-1},U_{it};\theta_0} = L_{f_{Y_{it}^+|X_{it},Y_{it-1},U_{it}}}^{-1} \left(\int \frac{\partial}{\partial \theta} f_{Y_{it}^+|X_{it},Y_{it-1},U_{it};\theta_0} f_{X_{it},Y_{it-1},X_{it-1},U_{it}} du_{it} \right). \quad (65)$$

These relationships can help expand the Kullback-Leibler function $K(\theta)$. Differentiating w.r.t. θ at $\theta = \theta_0$ leads to

$$\begin{aligned}
K'(\theta_0) &= \int \int \int \frac{\partial}{\partial \theta} \log(f_{X_{it}, Y_{it-1}, X_{it-1}; \theta}) \Big|_{\theta=\theta_0} f_{X_{it}, Y_{it-1}, X_{it-1}} dx_{it} dy_{it-1} dx_{it-1} \\
&= \int \int \int \frac{\frac{\partial}{\partial \theta} f_{X_{it}, Y_{it-1}, X_{it-1}; \theta}}{f_{X_{it}, Y_{it-1}, X_{it-1}; \theta}} \Big|_{\theta=\theta_0} f_{X_{it}, Y_{it-1}, X_{it-1}} dx_{it} dy_{it-1} dx_{it-1} \\
&= \int \int \int \frac{\partial}{\partial \theta} f_{X_{it}, Y_{it-1}, X_{it-1}; \theta} dx_{it} dy_{it-1} dx_{it-1} \\
&= \frac{\partial}{\partial \theta} \left(\int \int \int f_{X_{it}, Y_{it-1}, X_{it-1}; \theta} dx_{it} dy_{it-1} dx_{it-1} \right) \Big|_{\theta=\theta_0} \\
&= 0.
\end{aligned}$$

Thus, the true parameter θ_0 is a critical point of $K(\theta_0)$ and a sufficient condition of the uniqueness of θ_0 is $K''(\theta_0)$ is negative semidefinite. Denote E as the expectation w.r.t. $(x_{it}, y_{it-1}, x_{it-1})$. The second derivative of $K(\theta_0)$ in the scalar case is

$$\begin{aligned}
&K''(\theta_0) \\
&= \int \int \int \frac{\partial^2}{\partial \theta^2} f_{X_{it}, Y_{it-1}, X_{it-1}; \theta} \Big|_{\theta=\theta_0} dx_{it} dy_{it-1} dx_{it-1} \\
&\quad - \int \int \int \frac{\left(\frac{\partial}{\partial \theta} f_{X_{it}, Y_{it-1}, X_{it-1}; \theta} \right)^2}{f_{X_{it}, Y_{it-1}, X_{it-1}; \theta}} \Big|_{\theta=\theta_0} f_{X_{it}, Y_{it-1}, X_{it-1}} dx_{it} dy_{it-1} dx_{it-1} \\
&= \frac{\partial^2}{\partial \theta^2} \left(\int \int \int f_{X_{it}, Y_{it-1}, X_{it-1}; \theta} dx_{it} dy_{it-1} dx_{it-1} \right) \Big|_{\theta=\theta_0} - E \left[\left(\frac{\frac{\partial}{\partial \theta} f_{X_{it}, Y_{it-1}, X_{it-1}; \theta}}{f_{X_{it}, Y_{it-1}, X_{it-1}; \theta}} \Big|_{\theta=\theta_0} \right)^2 \right] \\
&= -E \left[\left(\frac{\frac{\partial}{\partial \theta} f_{X_{it}, Y_{it-1}, X_{it-1}; \theta}}{f_{X_{it}, Y_{it-1}, X_{it-1}; \theta}} \Big|_{\theta=\theta_0} \right)^2 \right] \tag{66}
\end{aligned}$$

where

$$\begin{aligned}
& \frac{\partial}{\partial \theta} f_{X_{it}, Y_{it-1}, X_{it-1}; \theta} \Big|_{\theta=\theta_0} \\
&= \frac{\partial}{\partial \theta} \frac{\int f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta} du_{it}}{\int \int \int \int f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta} du_{it} dx_{it} dy_{it-1} dx_{it-1}} \Big|_{\theta=\theta_0} \\
&= \int \underbrace{\frac{\partial}{\partial \theta} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}}_{\text{defined in Eq. (65)}} du_{it} \\
&\quad - \left(\int \int \int \int \underbrace{\frac{\partial}{\partial \theta} f_{X_{it}, Y_{it-1}, X_{it-1}, U_{it}; \theta_0}}_{\text{defined in Eq. (65)}} dx_{it} dy_{it-1} dx_{it-1} du_{it} \right) f_{X_{it}, Y_{it-1}, X_{it-1}}. \quad (67)
\end{aligned}$$

A similar derivation can be applied to the vector case and the form $K''(\theta_0)$, as Eq. (22) shows.

Q.E.D.

F. Proof of Lemma 3.3

Proof: First, suppose $\tilde{\mathcal{Y}}$ is a domain such that $\tilde{\mathcal{Y}} \subset \mathcal{R}$. Let the family $\{f(y|u) : u \in \mathcal{U}\}$ be complete in $L^2(\mathcal{R})$. For each $h \in L^2(\tilde{\mathcal{Y}})$ such that $\int_{\tilde{\mathcal{Y}}} h(y) f(y|u) dy = 0$ for all u . Extend h to a function in $L^2(\mathcal{R})$ by $\tilde{h}(x) = \begin{cases} h & \text{if } x \in \tilde{\mathcal{Y}}, \\ 0 & \text{otherwise.} \end{cases}$ It follows that $\int_{\mathcal{R}} \tilde{h}(y) f(y|u) dy = 0$ for all u . By the completeness of $f(y|u)$ over $L^2(\mathcal{R})$, $\tilde{h} = 0$. Thus, $h = 0$ and $f(y|u)$ is complete over $L^2(\mathcal{R})$. Thus, the completeness of a function over a smaller domain is implied by the completeness of the function over a larger domain, and sufficient conditions for the completeness of these two families can be reduced to the completeness in $L^2(\mathcal{R})$.

The family of functions $\{(y - c - u) \phi\left(\frac{y - c - u}{\sigma_\xi}\right) : u \in \mathcal{U}\}$ is complete in $L^2(\mathcal{R})$. Let $h(y) \in L^2(\mathcal{R})$ and $\int h(y) (y - c - u) \phi\left(\frac{y - c - u}{\sigma_\xi}\right) dy = 0$ for all $u \in \mathcal{U}$. Because $\frac{\partial}{\partial y} \phi\left(\frac{y - c - u}{\sigma_\xi}\right) = -\frac{y - c - u}{\sigma_\xi} \phi\left(\frac{y - c - u}{\sigma_\xi}\right)$, it follows that $\int h(y) \frac{\partial}{\partial y} \phi\left(\frac{y - c - u}{\sigma_\xi}\right) dy = 0$ for all $u \in \mathcal{U}$. Using the integration by part for each u leads to

$$\begin{aligned}
\int h(y) \frac{\partial}{\partial y} \phi\left(\frac{y - c - u}{\sigma_\xi}\right) dy &= h(y) \phi\left(\frac{y - c - u}{\sigma_\xi}\right) \Big|_{-\infty}^{\infty} - \int \frac{\partial}{\partial y} h(y) \phi\left(\frac{y - c - u}{\sigma_\xi}\right) dy \\
&= - \int \frac{\partial}{\partial y} h(y) \phi\left(\frac{y - c - u}{\sigma_\xi}\right) dy.
\end{aligned}$$

Applying the completeness of $\{\phi\left(\frac{y-c-u}{\sigma_\xi}\right) : u \in \mathcal{U}\}$ to this equation yields $\frac{\partial}{\partial y}h(y) = 0$, which implies that $h(y)$ is a constant function. The condition $h(y) \in L^2(\mathcal{R})$ makes $h(y) = 0$, proving the first completeness. As for the second completeness, suppose $h(y) \in L^2(\mathcal{R})$ such that $\int h(y) \left(\sigma_\xi^2 - (y - c - u)^2\right) \phi\left(\frac{y-c-u}{\sigma_\xi}\right) dy = 0$ for all $u \in \mathcal{U}$. Using $\frac{\partial^2}{\partial^2 y} \phi\left(\frac{y-c-u}{\sigma_\xi}\right) = -\frac{1}{\sigma_\xi^3} \left(\sigma_\xi^2 - (y - c - u)^2\right) \phi\left(\frac{y-c-u}{\sigma_\xi}\right)$ and the integration by part, rewrite the equation as

$$\begin{aligned} 0 &= \int h(y) \frac{\partial^2}{\partial^2 y} \phi\left(\frac{y-c-u}{\sigma_\xi}\right) dy \\ &= h(y) \frac{\partial}{\partial y} \phi\left(\frac{y-c-u}{\sigma_\xi}\right) \Big|_{-\infty}^{\infty} - \int \frac{\partial}{\partial y} h(y) \frac{\partial}{\partial y} \phi\left(\frac{y-c-u}{\sigma_\xi}\right) dy \\ &= -\frac{\partial}{\partial y} h(y) \phi\left(\frac{y-c-u}{\sigma_\xi}\right) \Big|_{-\infty}^{\infty} + \int \frac{\partial^2}{\partial^2 y} h(y) \phi\left(\frac{y-c-u}{\sigma_\xi}\right) dy \\ &= \int \frac{\partial^2}{\partial^2 y} h(y) \phi\left(\frac{y-c-u}{\sigma_\xi}\right) dy \end{aligned}$$

The completeness of $\{\phi\left(\frac{y-c-u}{\sigma_\xi}\right) : u \in \mathcal{U}\}$ implies that h satisfies the second order differential equation, $\frac{\partial^2}{\partial^2 y} h(y) = 0$. The characteristic equation of the differential equation is $r^2 = 0$. This suggests that the general solution of the differential equation is $h(y) = c_1 + c_2 y$, where c_1 and c_2 are constants. The condition $h(y) \in L^2(\mathcal{R})$ indicates $h(y) = 0$, reaching the second completeness. Q.E.D.

References

- AHN, S., AND P. SCHMIDT (1995): “Efficient Estimation of Models for Dynamic Panel Data,” *Journal of Econometrics*, 68(1), 5–28.
- AI, C., AND X. CHEN (2003): “Efficient Estimation of Models with Conditional Moment Restrictions Containing Unknown Functions,” *Econometrica*, 71(6), 1795–1843.
- ANDERSON, T., AND C. HSIAO (1982): “Formulation and Estimation of Dynamic Models Using Panel Data,” *Journal of Econometrics*, 18(1), 47–82.
- ANDREWS, D. (2011): “Examples of L^2 -Complete and Boundedly-Complete Distributions,” *Cowles Foundation for Research in Economics*.
- ARELLANO, M., AND S. BOND (1991): “Some Tests of Specification for Panel Data: Monte

- Carlo Evidence and an Application to Employment Equations,” *The Review of Economic Studies*, 58(2), 277–297.
- ARELLANO, M., AND O. BOVER (1995): “Another Look at the Instrumental Variable Estimation of Error Component Models,” *Journal of Econometrics*, 68(1), 29–51.
- BLUNDELL, R., AND S. BOND (1998): “Initial Conditions and Moment Restrictions in Dynamic Panel Data Models,” *Journal of Econometrics*, 87(1), 115–143.
- BOWDEN, R. (1973): “The Theory of Parametric Identification,” *Econometrica*, 41(6), 1069–1074.
- CANAY, I., A. SANTOS, AND A. SHAIKH (2011): “On the Testability of Identification in Some Nonparametric Models with Endogeneity,” *Working Paper*.
- CHEN, X., O. LINTON, AND I. VAN KEILEGOM (2003): “Estimation of Semiparametric Models When the Criterion Function is Not Smooth,” *Econometrica*, 71(5), 1591–1608.
- CHEN, X., AND X. SHEN (1998): “Sieve Extremum Estimates for Weakly Dependent Data,” *Econometrica*, 66(2), 289–314.
- CHERNOZHUKOV, V., I. FERNÁNDEZ-VAL, J. HAHN, AND W. NEWEY (2009): “Average and Quantile Effects in Nonseparable Panel Models,” *CeMMAP Working Paper*.
- D’HAULTFOEUILLE, X. (2011): “On the Completeness Condition in Nonparametric Instrumental Problems,” *Econometric Theory*, 1, 1–12.
- GRAHAM, B., AND J. POWELL (2008): “Identification and Estimation of ‘Irregular’ Correlated Random Coefficient Models,” Discussion paper, National Bureau of Economic Research.
- HAHN, J. (1999): “How Informative is the Initial Condition in the Dynamic Panel Model with Fixed Effects?,” *Journal of Econometrics*, 93(2), 309–326.
- HODERLEIN, S., AND H. WHITE (2009): “Nonparametric Identification in Nonseparable Panel Data Models With Generalized Fixed Effects,” *CeMMAP Working Paper*.
- HONORÉ, B. (1993): “Orthogonality Conditions for Tobit Models with Fixed Effects and Lagged Dependent Variables,” *Journal of Econometrics*, 59(1-2), 35–61.

- HONORÉ, B., AND L. HU (2004): “Estimation of Cross Sectional and Panel Data Censored Regression Models with Endogeneity,” *Journal of Econometrics*, 122(2), 293–316.
- HSIAO, C., M. HASHEM PESARAN, AND A. KAMIL TAHMISCIOGLU (2002): “Maximum Likelihood Estimation of Fixed Effects Dynamic Panel Data Models Covering Short Time Periods,” *Journal of Econometrics*, 109(1), 107–150.
- HU, L. (2002): “Estimation of a Censored Dynamic Panel Data Model,” *Econometrica*, 70(6), 2499–2517.
- HU, Y., AND S. SCHENNACH (2008): “Instrumental Variable Treatment of Nonclassical Measurement Error Models,” *Econometrica*, 76(1), 195–216.
- HU, Y., AND J. SHIU (2011a): “Identification of Semiparametric Measurement Error Models,” *Working Paper*.
- (2011b): “Nonparametric Identification Using Instrumental Variables: Sufficient Conditions For Completeness,” *Working Paper*.
- KIVIET, J. (1995): “On Bias, Inconsistency, and Efficiency of Various Estimators in Dynamic Panel Data Models,” *Journal of Econometrics*, 68(1), 53–78.
- NEWHEY, W., AND J. POWELL (2003): “Instrumental Variable Estimation of Nonparametric Models,” *Econometrica*, 71(5), 1565–1578.
- NEYMAN, J., AND E. SCOTT (1948): “Consistent Estimates Based on Partially Consistent Observations,” *Econometrica*, 93, 1–32.
- SHEN, X. (1997): “On Methods of Sieves and Penalization,” *Annals of Statistics*, 25, 2555–2591.
- SHIU, J., AND Y. HU (2010): “Identification and Estimation of Nonlinear Dynamic Panel Data Models with Unobserved Covariates,” *Economics Working Paper Archive*.
- WOOLDRIDGE, J. (2005): “Simple Solutions to the Initial Conditions Problem in Dynamic, Nonlinear Panel Data Models with Unobserved Heterogeneity,” *Journal of Applied Econometrics*, 20(1), 39–54.

Table 1: Simulation of Dynamic tobit model (N=250)

		Parameters			
	DGP	β_0	β_1	γ	σ_ξ^2
DGP I:	True value	0.2	-1	0	0.5
	Infeasible Mean	0.221	-1.004	-0.006	0.478
	Infeasible Median	0.234	-1.002	-0.005	0.470
	Standard error	0.139	0.116	0.108	0.086
	Mean	0.202	-1.009	-0.003	0.499
	Median	0.211	-1.006	-0.013	0.500
	Standard error	0.093	0.099	0.095	0.033
	Three-period Mean	0.201	-0.993	0.020	0.531
	Three-period Median	0.204	-0.995	0.018	0.544
	Standard error	0.097	0.109	0.104	0.105
DGP II:	True value	0.2	-1	0	0.5
	Infeasible Mean	0.204	-1.011	0.007	0.504
	Infeasible Median	0.190	-0.996	0.012	0.498
	Standard error	0.129	0.098	0.086	0.085
	Mean	0.185	-0.998	0.028	0.501
	Median	0.186	-0.985	0.033	0.503
	Standard error	0.084	0.101	0.094	0.034
	Three-period Mean	0.210	-1.018	0.010	0.552
	Three-period Median	0.210	-1.013	0.004	0.543
	Standard error	0.096	0.094	0.095	0.092
DGP III:	True value	0.2	-1	1	0.5
	Infeasible Mean	0.205	-1.012	1.008	0.501
	Infeasible Median	0.202	-1.009	1.004	0.497
	Standard error	0.113	0.087	0.080	0.057
	Mean	0.201	-1.059	0.876	0.518
	Median	0.196	-1.052	0.878	0.526
	Standard error	0.100	0.096	0.106	0.027
	Three-period Mean	0.210	-1.035	0.977	0.548
	Three-period Median	0.205	-1.036	0.982	0.544
	Standard error	0.095	0.107	0.122	0.038
DGP IV:	True value	0.2	-1	1	0.5
	Infeasible Mean	0.203	-1.011	1.008	0.505
	Infeasible Median	0.201	-1.005	0.997	0.504
	Standard error	0.111	0.090	0.079	0.058
	Mean	0.197	-1.062	0.881	0.548
	Median	0.197	-1.057	0.889	0.551
	Standard error	0.096	0.119	0.116	0.035
	Three-period Mean	0.191	-1.057	0.964	0.565
	Three-period Median	0.185	-1.071	0.964	0.557
	Standard error	0.103	0.126	0.146	0.075

Note: Standard errors of the parameters are based on the standard error of the estimates across 100 simulations. The three-period results are estimated using the sieve MLE in Shiu and Hu (2010).

Table 2: Simulation of Dynamic tobit model (N=500)

		Parameters			
	DGP	β_0	β_1	γ	σ_ξ^2
DGP I:	True value	0.2	-1	0	0.5
	Infeasible Mean	0.220	-1.017	0.007	0.505
	Infeasible Median	0.217	-1.018	0.018	0.504
	Standard error	0.123	0.075	0.063	0.074
	Mean	0.192	-1.015	-0.009	0.498
	Median	0.192	-1.018	-0.001	0.503
	Standard error	0.098	0.110	0.109	0.036
	Three-period Mean	0.207	-0.986	0.004	0.556
	Three-period Median	0.202	-0.988	-0.003	0.551
	Standard error	0.089	0.121	0.095	0.083
DGP II:	True value	0.2	-1	0	0.5
	Infeasible Mean	0.196	-1.001	0.004	0.500
	Infeasible Median	0.193	-1.005	0.004	0.497
	Standard error	0.091	0.077	0.069	0.056
	Mean	0.179	-0.991	-0.013	0.499
	Median	0.181	-0.995	-0.009	0.500
	Standard error	0.099	0.102	0.099	0.037
	Three-period Mean	0.197	-1.017	0.004	0.553
	Three-period Median	0.208	-1.026	0.014	0.554
	Standard error	0.097	0.105	0.100	0.033
DGP III:	True value	0.2	-1	1	0.5
	Infeasible Mean	0.193	-1.001	1.006	0.501
	Infeasible Median	0.196	-0.994	1.007	0.495
	Standard error	0.083	0.066	0.062	0.041
	Mean	0.196	-1.058	0.856	0.519
	Median	0.192	-1.058	0.860	0.519
	Standard error	0.105	0.090	0.093	0.028
	Three-period Mean	0.192	-1.047	0.976	0.564
	Three-period Median	0.193	-1.043	0.975	0.565
	Standard error	0.096	0.129	0.124	0.047
DGP IV:	True value	0.2	-1	1	0.5
	Infeasible Mean	0.201	-1.001	1.000	0.498
	Infeasible Median	0.210	-1.001	1.003	0.498
	Standard error	0.090	0.061	0.054	0.039
	Mean	0.186	-1.070	0.888	0.551
	Median	0.192	-1.072	0.899	0.554
	Standard error	0.090	0.123	0.103	0.036
	Three-period Mean	0.198	-1.068	0.937	0.573
	Three-period Median	0.198	-1.054	0.947	0.567
	Standard error	0.095	0.126	0.144	0.071

Note: Standard errors of the parameters are based on the standard error of the estimates across 100 simulations. The three-period results are estimated using the sieve MLE in Shiu and Hu (2010).

Table 3: Simulation of Dynamic tobit model (N=1000)

		Parameters			
	DGP	β_0	β_1	γ	σ_ξ^2
DGP I:	True value	0.2	-1	0	0.5
	Infeasible Mean	0.208	-1.006	0.005	0.490
	Infeasible Median	0.206	-1.005	0.001	0.489
	Standard error	0.063	0.049	0.040	0.043
	Mean	0.189	-0.967	-0.003	0.502
	Median	0.174	-0.973	-0.008	0.512
	Standard error	0.098	0.095	0.092	0.032
	Three-period Mean	0.208	-0.993	-0.005	0.550
	Three-period Median	0.216	-0.992	-0.004	0.554
	Standard error	0.093	0.112	0.107	0.044
DGP II:	True value	0.2	-1	0	0.5
	Infeasible Mean	0.210	-1.010	0.007	0.505
	Infeasible Median	0.202	-1.002	0.005	0.496
	Standard error	0.067	0.061	0.056	0.041
	Mean	0.199	-0.995	-0.006	0.503
	Median	0.200	-1.002	-0.008	0.509
	Standard error	0.107	0.102	0.105	0.509
	Three-period Mean	0.208	-0.996	-0.005	0.556
	Three-period Median	0.216	-0.999	-0.004	0.557
	Standard error	0.094	0.108	0.107	0.031
DGP III:	True value	0.2	-1	1	0.5
	Infeasible Mean	0.211	-1.006	1.005	0.499
	Infeasible Median	0.205	-1.004	1.005	0.498
	Standard error	0.058	0.052	0.050	0.027
	Mean	0.217	-1.051	0.839	0.526
	Median	0.230	-1.053	0.836	0.528
	Standard error	0.087	0.101	0.096	0.026
	Three-period Mean	0.205	-1.014	0.952	0.568
	Three-period Median	0.206	-1.014	0.975	0.566
	Standard error	0.092	0.114	0.136	0.047
DGP IV:	True value	0.2	-1	1	0.5
	Infeasible Mean	0.208	-1.004	1.004	0.499
	Infeasible Median	0.208	-1.008	1.005	0.497
	Standard error	0.058	0.049	0.046	0.026
	Mean	0.204	-1.058	0.879	0.558
	Median	0.213	-1.063	0.880	0.559
	Standard error	0.090	0.116	0.101	0.032
	Three-period Mean	0.210	-1.067	0.917	0.590
	Three-period Median	0.213	-1.052	0.930	0.577
	Standard error	0.099	0.133	0.144	0.068

Note: Standard errors of the parameters are based on the standard error of the estimates across 100 simulations. The three-period results are estimated using the sieve MLE in Shiu and Hu (2010).

Table 4: Simulation of Average Partial Effects

		State Dependence $SD(\bar{X}_{it}, \bar{Y}_{it-1})$		
DGP		N=250	N=500	N=1000
DGP I:	True value	0	0	0
	Mean	0.002	-0.002	-0.001
	Median	0.002	-0.003	0.001
	Standard error	0.008	0.008	0.009
DGP II:	True value	0	0	0
	Mean	-0.001	-0.001	0.001
	Median	0.002	0.001	0.002
	Standard error	0.008	0.010	0.010
DGP III:	True value	0.579	0.579	0.579
	Mean	0.660	0.652	0.644
	Median	0.665	0.655	0.652
	Standard error	0.115	0.093	0.082
DGP IV:	True value	0.479	0.479	0.479
	Mean	0.618	0.620	0.646
	Median	0.632	0.635	0.645
	Standard error	0.177	0.139	0.145

Note: Standard errors of the parameters are based on the standard error of the estimates across 100 simulations. $SD(\bar{X}_{it}, \bar{Y}_{it-1}) \equiv \mu_1(\bar{X}_{it}, \bar{Y}_{it-1}) - \mu_1(\bar{X}_{it}, 0)$, where $(\bar{X}_{it}, \bar{Y}_{it-1})$ is the mean of (X_{it}, Y_{it}) and it is the difference of the average structural functions of two different outcomes of Y_{it-1} , 0 and \bar{Y}_{it-1} . This represents the magnitude of the state dependence.

Table 5: Sample Statistics

Variable	Definition	Period _{it}	Period _{t+1}
Lnexp	log(medical expenditures+1)	5.292 (2.903)	5.307 (3.038)
Lninc	ln(family income+1)	9.056 (2.821)	9.217 (2.695)
Lnfam	ln(family size)	1.036 (0.538)	1.034 (0.542)
Age	Age	39.427 (12.498)	40.429 (12.500)
Male	=1 if person is male; 0 otherwise	0.469 (0.499)	0.469 (0.499)
Black	=1 if race of household head is black; 0 otherwise	0.148 (0.355)	0.148 (0.355)
Education	Education of the household head	12.599 (3.087)	12.599 (3.087)
Physical	=1 if the person has a physical limitation; 0 otherwise	0.057 (0.231)	0.059 (0.235)
Ndental	Number of dental care visits	0.938 (1.746)	0.857 (1.617)
Good	=1 if self-rated health is good; 0 otherwise	0.266 (0.442)	0.276 (0.447)
Fair	=1 if self-rated health is fair; 0 otherwise	0.086 (0.280)	0.081 (0.274)
Poor	=1 if self-rated health is poor; 0 otherwise	0.026 (0.158)	0.027 (0.162)
Deduction	=1 if the person has nonzero itemized deductions; 0 otherwise	0.057 (0.232)	0.054 (0.227)
Medicare	=1 if the person is covered by Medicare; 0 otherwise	0.025 (0.156)	0.034 (0.182)
Medicaid	=1 if the person is covered by Medicaid; 0 otherwise	0.070 (0.255)	0.068 (0.253)
Sample size		7,669	7,669

Note: The variables in Period_{it} and Period_{t+1} refer to the first-year and the second-year values of each participation respectively. There are 1,430 and 1,578 individuals with zero medical expenditures in Period_{it} and Period_{t+1}, respectively. Standard deviations are in parentheses.

Table 6: Panel Censored Estimates for Health Expenditure

	Linear Fixed Effects (1)	RE Tobit (2)	Semi-parametric Dynamic Tobit (3)		
	Coefficient	Coefficient	APE	Coefficient	APE
Lnexp _{it-1}	— —	— —	— —	1.052*** (0.001)	1.448*** (0.006)
Lninc	0.031*** (0.008)	0.042*** (0.011)	0.039*** (0.001)	0.041*** (0.001)	0.056*** (0.001)
Lnfam	-0.252*** (0.045)	-0.299*** (0.056)	-0.276*** (0.003)	-0.301*** (0.001)	-0.414*** (0.002)
Age	0.040*** (0.002)	0.048*** (0.003)	0.044*** (0.001)	0.050*** (0.001)	0.068*** (0.001)
Male	-1.130*** (0.049)	-1.399*** (0.062)	-1.294*** (0.032)	-1.399*** (0.001)	-1.927*** (0.008)
Black	-0.581*** (0.069)	-0.717*** (0.086)	-0.653*** (0.012)	-0.717*** (0.001)	-0.987*** (0.004)
Education	0.145*** (0.008)	0.184*** (0.011)	0.170*** (0.002)	0.181*** (0.001)	0.250*** (0.001)
Physical	0.806*** (0.098)	0.854*** (0.119)	0.788*** (0.007)	0.855*** (0.001)	1.177*** (0.005)
Ndental	0.442*** (0.012)	0.496*** (0.015)	0.458*** (0.004)	0.502*** (0.001)	0.691*** (0.003)
Good	0.342*** (0.047)	0.391*** (0.059)	0.362*** (0.008)	0.392*** (0.001)	0.539*** (0.002)
Fair	1.037*** (0.080)	1.180*** (0.098)	1.111*** (0.031)	1.178*** (0.001)	1.621*** (0.007)
Poor	1.777*** (0.142)	1.956*** (0.173)	1.865*** (0.054)	1.957*** (0.001)	2.694*** (0.011)
Deduction	0.384*** (0.089)	0.432*** (0.108)	0.402*** (0.010)	0.429*** (0.001)	0.590*** (0.003)
Medicare	0.900*** (0.143)	0.995*** (0.175)	0.936*** (0.035)	0.995*** (0.001)	1.370*** (0.006)
Medicaid	1.138*** (0.092)	1.346*** (0.114)	1.270*** (0.001)	1.343*** (0.001)	1.849*** (0.008)

Note: Bootstrap (simulation) standard errors are reported in parentheses, using 100 bootstrap replications. APEs are reported by taking derivatives or differences of ASF at the sample mean of (x_{it}, y_{it-1}) .