

**Health Human Capital in Developing Countries:
Evidence from Infant Mortality Rates and Adult Heights**

Harvard Center for Population and Development Studies

Harvard School of Public Health

Yoko Akachi^a and David Canning^a

October 2009

Abstract

We investigate trends in health human capital using data on infant mortality rates and adult heights for birth cohorts from 1960 to 1985 from 39 developing countries. In Sub-Saharan Africa, despite declining infant mortality rates, adult heights have not risen. We argue that childhood exposure to disease, and nutrition, have remained stagnant in Sub-Saharan Africa and the decline in infant mortality has been due to technological progress in preventing the death of children in poor health rather than improved morbidity. Despite falling infant mortality rates, the health human capital and worker productivity of adults in Sub-Saharan Africa may remain low.

Keywords: adult height, stature, human capital, Sub-Saharan Africa, disease burden

Corresponding Author:

David Canning

E-mail: dcanning@hsph.harvard.edu

Program on the Global Demography of Aging

Harvard Center for Population and Development Studies

9 Bow Street

Cambridge, MA 02138

Tel: (617)432-6336

Fax: (617)495-5418

The authors gratefully acknowledge funding from the National Institute on Aging, Grant No. 1 P30 AG024409-01.

1. Introduction

Over the last fifty years the world has seen enormous improvements in population health in terms of falling mortality rates. These reductions in mortality rates, and gains in life span, have created large improvements in human welfare (Becker, Philipson, and Soares 2005). Improvements in mortality in Europe and North America in the 19th century were associated with improved nutrition and public health measures, such as the provision of clean water and sanitation (Cutler, Deaton, and Lleras-Muney 2006, McKeown 1983). These mortality improvements went hand in hand with reductions in childhood morbidity and improved childhood nutrition, leading to increases in adult stature (Fogel and Costa 1997, Alter 2004, Crimmins and Finch 2006).

In addition to the direct welfare gains from improved health we can also view health as a form of human capital. Healthier workers are more productive. It may be, however, health investments in children rather than investments in adults that are most important for human capital. There is increasing evidence that childhood health and nutrition can have a substantial impact on both physical and cognitive development, and eventual health status and productivity as an adult (Mendez and Adair 1999, Elo and Preston 1992, Hayward and Gorman 2004, O'Rand and Hamil-Luker 2005, Barker et al 1989, Kuh 2002, Yi, Gu, and Land 2007, Hodinott et al 2008). In addition to increasing worker productivity, improved health can also have more indirect effects on the economy; longer life spans can change life cycle behavior and the incentive to investment in human and physical capital. Reduced mortality will also influence population growth and age structure which may have effects on aggregate economic outcomes (Bloom and Canning 2000).

While there is increasing evidence that health is an important component of human capital, there is a mismatch in how health is measured at the individual and population levels. At

the individual level we can use measures of individual well-being and morbidity, such as self reported health, anthropometrics, medical records, biomarkers for the presence of disease, or tests of specific abilities (Schultz 2005). At the aggregate level, however, we lack databases that provide summary statistics for these measures for populations. In cross country studies researchers usually use mortality rates (or some constructed indicator based on them, such as life expectancy) as their measure of population health (Bloom, Canning, and Sevilla 2004). It is clear that worker productivity is related to well-being and morbidity of living workers, not to mortality rates; nevertheless in macroeconomic studies, mortality indicators such as life expectancy are most often used as a proxy for the health of the living population.

This raises the question of whether the mortality measures used at the population level are related to the morbidity measures used at the individual level. Weil (2007) identifies this as a central issue in comparing household level and macroeconomic studies of the effect of health on the economy. He argues that if we assume that health is a one-dimensional construct, different measures of health should be closely related. He shows that in a small set of currently developed countries, for which a long time span of data is available, life expectancy and average adult height tend to move together over time. Individual level studies have found a strong connection between variations in adult height (due to differences in childhood health and nutrition) and wages (Schultz 2002, Schultz 2005). Weil uses the link between long run changes in height and changes in life expectancy, and the estimated effect of height on productivity, to calibrate the effect of improvements in life expectancy on the economy.

We extend this inquiry to see if the link between reductions in mortality rates and improvements in adult height holds in developing countries over the last 50 years. We focus on infant mortality rather than life expectancy as our mortality measure. This is partially because it is child health that is central to adult well-being. It is nonetheless also the case that for most

developing countries we lack comprehensive data on age specific mortality rates, apart from infant mortality rates, and life expectancy figures are based on mortality rates extrapolated from infant and child mortality using model life tables. This means that in practice, in many developing countries, reported life expectancy is a transformation of infant and child mortality (Deaton 2006).

We think of infant mortality and adult height as depending on nutrition and the burden of disease experienced in childhood. While we have a measure on nutrition, we think of disease burden as a latent variable that affects both a cohort's infant mortality and its eventual adult height. In our theory section below we set out a model where cohort infant mortality and eventual adult height both depend on childhood nutrition and disease burden and show how this implies that a cohort's experience of infant mortality and average adult height should be linked.

We use from data for 39 developing countries, for cohorts born from 1961 onwards. As in Weil (2007) we find little connection between mortality rates and adult heights across countries. Adult Africans are tall compared to people in other countries, despite poor nutrition and high mortality rates when young. We argue that these cross country differences are due to unobserved fixed factors, not child health. These country fixed effects could to some extent reflect genetic differences between people in different countries but other factors, such as selection effects, could also play a role (Deaton 2007).

When we examine trends in cohort infant mortality rates, and average adult height, over time, within countries, we find a strong correlation; the countries with the greatest improvements in infant mortality are also those with the largest increases in average adult height. We ascribe this common movement to changes in underlying child health. While all countries in our sample have seen falling infant mortality rates, we find that in Sub Saharan Africa as a whole adult heights have not been increasing, and in some countries we even see significant declines in

height.

Most developed countries underwent a transition in which infant mortality fell and adult height increased. We argue that this pattern continues to hold true today in much of Latin America and Asia where reductions in infant mortality go hand in hand with improved nutrition, reductions in child morbidity, and increases in adult heights. In Sub-Saharan Africa, however, we find a distinctively different pattern: while infant mortality has been falling, adult heights have been stagnating, or even declining, over the last 50 years. This undermines the view that since infant mortality rates have been falling health human capital has been rising. There has been no improvement in health human capital, as measured by height, in the region.

One possible reason behind the divergent trends in infant mortality and adult height in Sub-Saharan Africa is the source of the infant mortality reductions in the region. In most other regions of the world, nutrition, as measured by energy (calorie) or protein intake per person, has been increasing. Sub-Saharan Africa in contrast has experienced little in the way of increases in nutrition. Rather than broad based improvements in nutrition and public health measures such as access to clean water and sanitation, mortality reduction in Sub-Saharan Africa appears to have occurred through health interventions measures that directly reduce mortality with limited effect on disease prevalence, morbidity, and the physical development of children.

The idea that reductions in infant mortality in Africa have not been associated with improved health in surviving children has been suggested before. Huffman and Steel (1995) and Guerrant, Carneiro-Filho, and Dillingham (2003) argue that oral rehydration therapy and measles vaccination have large effects on infant mortality rates, while morbidity rates remain high and the physical development of children continues to be impaired. Greenwood (1987) finds little improvement in the stunting and wasting of children in rural Gambia, despite high levels of vaccination coverage, and few deaths due to vaccine preventable diseases. Similarly, Pinchinat et

al. (2004) find that vaccination and malaria prevention in Senegal improve childhood survival but do not improve the health status of living children. These researchers attribute the persistent high child morbidity rates, despite improvements in child mortality, to continuing acute respiratory infections, malaria, chronic diarrhea, and malnutrition.

Our research suggests that health is multi-dimensional, and that changes in infant mortality rates and adult heights, while tending to move together overall, may diverge significantly in some regions. Improvements in infant mortality in Africa may not be indicative of broad based improvements in population health and health human capital.

2. Height and Infant Mortality Rates as Health Indicators

Cohort heights have been used by economic historians (Fogel 1993, Komlos 1993, Steckel 1995) as measures of “biological standard of living” when other indicators have not been available. It has been suggested that height can also be used as an indicator of health status in modern populations (Komlos and Lauderdale 2007). Steckel (2009) provides a recent survey of work in the social sciences using height as an indicator while Steckel (2008) discusses heights and mortality rates as measures of well being. Several studies have investigated the link between childhood health, nutrition, and adult height at the population level. Across developing countries, greater average protein intake is associated with greater average adult height (Jamison, Leslie, and Musgrove 2003). In addition, recent data has been used to examine the determinants of adult height in Sub-Saharan Africa (Moradi 2002, Moradi 2006, Deaton 2007, Akachi and Canning 2007).

Physical growth during childhood and adolescence, and the adult height that it ultimately leads to, is an indicator of childhood nutrition and the disease environment. Although height has

a genetic component¹, it is also significantly affected by childhood living conditions. The major proximate determinants of height are nutrition, disease environment, and work intensity (Tanner 1992, Peck and Lundberg 1995, Steckel 1995, Brush, Harrison, and Waterlow 1997, Stephensen 1999). The disease environment matters for physical development because infection can reduce appetite, and cause diarrhea and other health conditions that prevent absorption of nutrients and physical growth. The more distal socioeconomic determinants of height include income, inequality, public health measures such as water and sanitation, access to health care, personal hygiene, health technologies, labor organization, cultural values, and food prices (Silventoinen 2003).

We assume that adult heights depend on nutrition intake and the morbidity experienced in childhood due to the disease environment.² More formally, let us suppose average height, h_{it} , in country i at time of the cohort born at time t depends on a vector of variables x_{it} . This vector includes genetic endowment and other environmental conditions when the cohort is young. We single out childhood nutrition n_{it} and childhood disease burden, d_{it} , as factors influencing height. We then assume that height of the cohort born in year t in country i is determined according to the equation

$$h_{it} = \alpha_x x_{it} + \alpha_n n_{it} + \alpha_d d_{it} + \varepsilon_{it} \quad (1)$$

¹ At the individual level there is evidence from developed countries that as much as 80% of variation in the heights of individuals can be ascribed to genetic factors (Silventoinen 2003, Visscher 2006). Up to 20% of variation in body height is thus attributed to environmental variation in Western modern societies (Silventoinen 2000, Stunkard 1986). In poorer environments, this proportion is likely to be larger, with lower heritability of body height as well as larger social economic body height differences.

² The net nutrition approach would also include the effect of labor and other physical activity in childhood that consumes energy and reduces the remaining energy balance available for physical growth.

where ε_{it} is an error term. We assume that the infant mortality rate in country i at time t also depends on the same exogenous factors (if the factors affecting height and infant mortality differ they can be included in the other equation with a coefficient of zero) and childhood health according to:

$$m_{it} = \beta_x x_{it} + \beta_n n_{it} + \beta_d d_{it} + u_{it} \quad (2)$$

where u_{it} is an error term. These equations can be regarded as a linearized version of the model of height and mortality proposed by Alter (2004). The difficulty with equations (1) and (2) is that childhood disease burden d_{it} is an unobserved latent variable. We can combine equations (1) and (2) by substituting for the latent variable d_{it} to give the relationship

$$h_{it} = \left(\alpha_x - \frac{\alpha_d}{\beta_d} \beta_x \right) x_{it} + \left(\alpha_n - \frac{\alpha_d}{\beta_d} \beta_n \right) n_{it} + \left(\frac{\alpha_d}{\beta_d} \right) m_{it} + \left(\varepsilon_{it} - \frac{\alpha_d}{\beta_d} u_{it} \right) \quad (3)$$

We see that the height and infant mortality rate of a cohort will be linked because both are linked to the latent variable, childhood disease burden. Akachi and Canning (2007) estimate the model given by (3) using data from Sub-Saharan Africa and find a significant relationship between the infant mortality rate in childhood and adult height even after controlling for income and nutrition.

However equation (3) does not allow us to estimate the underlying disease burden. To do this we replace the unobserved latent variable in equations (1) and (2) with a country specific intercept and time trend specified by

$$d_{it} = d_{i0} + d_i t \quad (4)$$

This means that equations (1) and (2) can be written as

$$h_{it} = \alpha_x x_{it} + \alpha_n n_{it} + \alpha_d d_{i0} + (\alpha_d d_i) t + \varepsilon_{it} \quad (5)$$

$$m_{it} = \beta_x x_{it} + \beta_n n_{it} + \beta_d d_{i0} + (\beta_d d_i) t + u_{it} \quad (6)$$

We estimate the evolution of the childhood disease burden as a country specific fixed effect and

linear trend. We assign the fixed effect and trend in height not explained by other variables to the disease burden. Similarly we explain the fixed effect and time trend in infant mortality not explained by other factors to this same latent variable. In principle, equations (5) and (6) can give us two independent estimates of the level and time trend in the disease burden. We produce these estimates below and compare the results from the two regressions. If our model is correct, and we control properly for other factors affecting height and infant mortality, the estimates of the levels and trends in disease burden for each country produced by equations (5) and (6) using data on adult heights and infant mortality respectively should be very similar.

An important question is what additional variables to add to the model in the vector x_{it} that may help explain infant mortality and adult height. We begin with estimates that do not include any additional variables, even nutrition. In this case we are essentially assuming that child health is the only factor influencing infant mortality height. In this case, unadjusted data on infant mortality rates and adult height should display the same pattern over time and across countries. Omitted variables in the model will mean that estimates of child health from height and infant mortality do not agree. In this case the fixed effects and time trends in height and infant mortality will reflect these omitted variables. We investigate what variables are required as additions to the model to bring the implied measures of child health, and disease burden into agreement.

One possible set omitted of variables are genetic variations in population height. These genetic variations may produce differences in height across populations. However, we assume that these variations are fixed over the short time period we consider so that they may affect the country fixed effects in equation (5) but not the country specific time trends. A second potential omitted variable is technological progress in mortality reduction. At each level of child health we may see reductions in mortality over time due to the introduction of new life saving interventions.

If these technologies save lives, but leave the underlying health condition and morbidity unchanged, we will see a time trend in equation (6) that is not reflected in heights. Both these potential omitted variables will reduce the correlation between heights and infant mortality rates unless we control for them.

We also estimate equations (5) and (6) adding average protein consumption, and average calorie consumption, in the cohort's year of birth as explanatory variables that proxy the cohort's food consumption. This removes variations in height and infant mortality due to these nutrition factors and we can see if the remaining variations in height and infant mortality, due to disease burden, move together. This approach mirrors Alter (2004) who thinks of food consumption and disease environment as the two factors influencing height and infant mortality. We can regard our estimates of child health for our unadjusted model as combining the effects of nutrition and the disease environment. Once we adjust for nutrition our latent variable is a measure of the disease environment.

3. Data

The height data we use are for women and come mainly from Demographic and Health Surveys (DHS), though we also use Family Life Surveys (FLS) for Mexico and Indonesia. All Demographic and Health Surveys available at the time of analysis that include female height as a variable were employed in this analysis (with the exception of Egypt, for which we found some inconsistency in the data, see below). Not all countries with DHS have adult height data. Height in Demographic and Health Surveys is measured by the interviewer, using a headboard. The typical Demographic and Health Surveys dataset contains the heights of women from age 15 to 49 for a nationally representative sample. We use the sampling weights provided to construct an

average height for cohorts by birth year. We only use heights of women aged 20 and above on the grounds that at age 20, physical growth has usually ceased.

One complication is that in earlier surveys, only the heights of mothers with children under age 5 were measured, while in later Demographic and Health Surveys the height of all women 15 to 49 was measured. This creates a sample selection problem since mothers are not a random sample. For example, if higher socioeconomic status is associated with fewer children, and height is positively linked to socioeconomic status, then the average height of mothers may be lower. In our check on data consistency, we examined the average height of each cohort as measured in different DHS surveys and found them to be remarkably similar, independently of whether the data were for all women or just mothers. These findings agree with those of Moradi (2006), who argues that there is little selection bias due to this issue in developing countries because the vast majority of adult women have children.

Table 1 shows, as examples, some descriptive statistics for the distribution of cohort heights for cohorts born in 1960, 1965, 1970 and 1975 from the Bangladesh 2004, Bolivia 2003, and Ghana 2003 DHS. The standard deviation of individuals' heights is around 6 centimeters; this distribution of heights is fairly uniform in all the samples we use. There is some evidence of a positive (right) skew in the distribution of recent cohorts for Bangladesh and Bolivia DHS while there appears to be a negative skew for the distribution in Ghana. Positive kurtosis (so that the peak of the distribution is higher and narrower, with fatter tails, than the normal) was observed for each cohort for all three country DHS. Tests of the hypothesis that the distribution is normal fails to be rejected for the three 1960 cohorts and 1970 Ghana cohort, but it is rejected for the other cohorts. Figure 1 shows the estimated distribution of heights of the cohort born in 1970 in Ghana 2003 DHS. The distributions are also shown for the 1975 cohort in Bolivia 2003 DHS (Figure 2).

The rejection of normality was common in many of our datasets. The deviation from normality could be taken as evidence of a selection effect, which could potentially be corrected by statistical methods. However, it is possible that differential health and nutrition across individuals creates a non-normal distribution and that a selection "correction" would bias the results (e.g. see Jacobs, Katzur, and Tassenaar 2004).

We construct average height for each cohort by year of birth from each survey. For countries with multiple DHS surveys, cohort heights by birth years were graphed to check for consistency when the same cohort is included in different surveys. The example for the three DHS surveys of Ghana is shown in Figure 3, and that of Bolivia in Figure 4. In most cases the results for average height by birth cohort using different surveys were very similar. The only country in which we found considerable variation was Egypt. Egypt has three DHS surveys, one of which, the 2003 survey gives heights that consistently differ from the 1995 and 2000 surveys (Figure 5). We are uncertain of the reason for these differences and the surveys from Egypt were discarded.

In order to construct figures for average cohort height we average over available DHS surveys for each country when multiple surveys are available. This gives larger sample sizes except at the beginning and end of the series where often only data from one survey is available. Larger sample size reduces the measurement error in the cohort average and result in noticeably lower volatility in average heights. The number of observations for each cohort from a survey is used as a weight in taking the averages for a cohort across different surveys.³ Countries with a single DHS survey give about 30 years of cohort data (women aged 20 to 49 are sampled to create cohort heights). For countries with multiple DHS surveys we get longer time spans of

³ We calculate the average height of a cohort from each survey using its sampling weights and then combine averages for the same cohort from different surveys weighting by the sample size of the cohort in the survey.

cohort height data though most of the information gained from using additional surveys is in additional observations of women's height from the same cohort.

We compare heights with indicators for infant mortality and nutrition. We use the infant mortality rate from the World Bank's World Development Indicators (2005), with data going back to 1960. We interpolated the infant mortality rate over gaps of up to two years to derive an annual time series. For nutrition we use daily average consumption of calories and protein from the World Food Organization (2006) FAOSTAT database, with data going back to 1961. The Food and Agriculture Organization (FAO) food balance sheets calculate the consumption of each foodstuff. Consumption of foodstuffs is calculated from two sources: national food supplies given by production plus imports, minus exports and wastage, and data on food consumption from household surveys. Jacobs and Sumner (2002), discuss the construction of the food balance sheets, problems in constructing the data, and their appropriate use. Calories and Protein consumed per capita are calculated from national consumption of each foodstuff using nutritional tables of calorie and protein content, and dividing by the population. These estimates for national food or nutrient availability do not account for the distribution of food or nutrient supply between regions within a country or among other groups of households. The data we use provide daily average consumption of calories and protein for each country by year from 1961 to 2002.

We begin by investigating the link between physical development, infant mortality, and nutrition, looking at how these change over time in developing countries. Table 2 shows the time trends for infant mortality rates, protein, and calorie intake for countries in Sub-Saharan Africa, from 1961 to 1985, matched with the trends in adult height for cohorts born during that period. In every country in the region except Rwanda, we see statistically significant declines in the infant mortality rate. In terms of protein and calorie intake, the picture in Sub-Saharan Africa is much more mixed: as many countries have seen declines in nutritional intake as have seen

increases. Finally, adult heights have risen significantly in only two countries, Kenya and Senegal, whereas three countries, Chad, Ethiopia and Rwanda, have seen heights decrease.

We can compare these trends with those in developing countries outside Sub-Saharan Africa over the same time period, shown in Table 3. Infant mortality rates fell significantly in every country outside Sub-Saharan Africa. Nutrition in the form of either calorie or protein intake increased significantly in every country except Bangladesh, Nicaragua, and Peru. Adult heights also increased significantly in every country except Bangladesh, where they stagnated.

These national trends are summarized in Table 4 where we report regional time trends that represent the average annual change in the variable in the region.⁴ In terms of infant mortality, we find similar rates of decline in Sub-Saharan Africa and developing countries in other regions: a decrease of about 1.7 versus 2.4 infant deaths per thousand live births each year.⁵ On the other hand, while both protein and calorie consumptions have been increasing significantly elsewhere, within Sub-Saharan Africa protein and calorie consumption remained virtually unchanged over the whole period. The trends in height are also quite distinct. In Sub-Saharan Africa, heights overall have been decreasing; the cohort born in 1985 is about 0.5 centimeters shorter than the cohort born in 1961. In contrast, in the rest of the developing world, the height of adult women has risen by approximately 1.8 centimeters on average during this twenty-four year period.

4. Results

We begin by estimating equation (3). If our theory is correct there should be a relationship

⁴ We average over countries treating each as an observation rather than construct population weighted averages.

⁵ This estimate is based on 1961-1985. Using the same dataset, Adetunji and Bos (2006) finds that in Sub-Saharan Africa as a whole, infant mortality rates declined from 149 per 1,000 live births in the 1960s to about 101 in 2005, or annual change about 1.2, about half our estimate. The difference in the estimates is likely due to the HIV/AIDS epidemic that affected the regions since the 1990's during which the decline in infant mortality stagnated.

between the infant mortality rate and adult height, after controlling for other variables. Results estimating this link are given in Table 5. In column 1 of Table 5 we simply estimate the effect of the infant mortality rate in a cohort's year of birth on the height of the cohort. Estimation is by weighted least squares where we weight each observation by the number of observations used to calculate the cohort's average height. A larger number of observations reduces the random sampling error in estimated cohort height and gives a more reliable figure. We find a positive coefficient on infant mortality indicating that cohorts that experience a higher infant mortality rate when young are taller as adults. This corresponds to the fact that Sub-Saharan Africa has the highest infant mortality rates, but also the tallest adults (Deaton 2007).

In column 2 of Table 5 we re-estimate the relationship with country fixed effects. We now find a significant negative coefficient on infant mortality. Within countries, cohorts that experience low infant mortality rates as children are taller as adults. The country fixed effect may reflect genetic or environmental sources of variation in height across countries (Ruff 2002). At this point, there is little we can infer from the fixed effects and what they may mean. In principle, they represent country specific, time invariant, omitted variables that affect height. They may reflect genetic differences between the populations, but genetic effects alone may unlikely be the sole cause of these significant fixed effects. There is also the omitted variable problem; the country fixed effects could capture forces such as inequality in nutrition, which are omitted from the model.

In column 3 we add calorie and protein consumption per capita in the cohort's year of birth as control variables. We find that within countries the tallest cohorts are those that had low infant mortality and high levels of protein consumption in childhood. There has been some discussion on whether protein or calorie intake matters more to physical growth. Hegsted (1971) puts more weight on calorie intake, but protein is often considered as the most important single

nutrient affecting growth (Zerfas 1986, Allen 1994, Martorell & Habicht 1986). Martorell (1976) argues that the relative contribution of calories and proteins to the association depends upon which nutrient is limiting in the current diet.

In column 4 of Table 5 we include a worldwide time trend to allow for technical progress that might change the relationship between the disease environment and infant mortality. We find a significant negative time trend in heights though the significant effect of childhood infant mortality and protein remain. We do not think there is really a downward trend in heights over time. Rather as we shall see below there appears to be time trends in infant mortality that are not reflected in heights. The results in Table 5 suggest a relationship between infant mortality and height occurs only after we control for country fixed effects. The results on the relationship between infant mortality and adult height are similar to those in Akachi and Canning (2007) who focus only on Africa but allow for the effects of infant mortality and nutrition at different points in childhood and not just in the year of birth. .

We now turn to estimates of child health and disease based on our data on infant mortality and heights. We begin with the model set out in equations (5) and (6) without any additional controls. That is, we regress cohort height by year of birth on country fixed effects and time trends and then do the same for the infant mortality rate. This gives us an estimated initial level and time trend for each variable in each country. We do not report the initial levels for each country but the time trends in height and infant mortality are as reported in Table 2. Figure 6 plots the initial levels of infant mortality against the initial level of height, the country specific fixed effects, for each country from this regression. There is no evidence of a relationship. However we do see that at each level of infant mortality people in sub-Saharan Africa appear to be taller. Figure 6 suggests that if we take the level of infant mortality, and population average heights, as our two measures of population health, the two measures do not agree.

In Figure 7 we plot the time trend in infant mortality against the time trend in height for each country. We think of a positive time trend in height as evidence of increasing child health. We think of a negative time trend in infant mortality also as evidence of increasing child health. We see a strong relationship in this graph with countries that have the largest decreases in infant mortality experiencing the largest gains in height. Both Sub-Saharan African countries and non Sub-Saharan African countries appear to follow the same relationship in trends. Nonetheless we see that Sub-Saharan African countries tend to have smaller declines in infant mortality than in other regions and it tends to have decreasing rather increasing heights. Adult heights and infant mortality rates are telling us the same story about child health in terms of the ranking of countries. Outside Sub-Saharan Africa, both indicators provide evidence of improvements in child health. In Sub-Saharan Africa, however, falling infant mortality rates indicate improving child health while decreasing adult heights suggest declining child health.

The difference in implied child health from the two measures, both in levels in changes, indicates that we have unobserved variables that are affecting the outcomes. We therefore estimate equations (5) and (6) adjusting for covariates x_{it} . The country fixed effects, and time trends in (5) and (6), after adjusting for these covariates, should tell the same story.

We begin in Table 6 by estimating equation (5) for adult height. In column 1 of Table 5 we control for calorie and protein consumption per capita in the year of birth. While protein consumption appears to have a positive effect on heights, the coefficient on calorie consumption is negative and significant. However, column 1 does not control for country fixed effects – the result is due to the fact that in Sub-Saharan Africa people are tall despite having very low calorie consumption in childhood. In column 2, controlling for country fixed-effects, we find a positive effect of calorie consumption while protein consumption is not statistically significant. This seems more reasonable and suggests that when childhood calorie consumption rises over

time later cohorts are taller.

In column 3 of Table 6 we add a worldwide time trend, but this is not significant. Once we control for nutrition there is no trend in heights. In column 4 of Table 6 we add country specific time trends. These make both our nutrition variables insignificant. The long run trend in calorie consumption in a country explain the long run trend in height in that country. However, short run movements in calorie consumption do not explain movements around the trend. Measurement error in our nutrition data may make it a good indicator of trends in the long run but a poor indicator of year to year changes.

In Table 7 we repeat the analysis in table 6 but with the log infant mortality rate as the dependent variable. In columns 1 and 2, with and without fixed effects respectively, we find conflicting evidence on the effect of nutrition with higher calorie consumption appearing to lower infant mortality rates while higher protein consumption appears to raise them. However in column 3, where we add a world wide time trend, improvements in nutrition in either form lower infant mortality. Column 3 of Table 7 indicates that there is a long run trend towards lower infant mortality, with mortality rates falling about 1.8% a year across all countries independently of changes in nutrition. We think of this effect as technological progress in health care. Cutler, Deaton, and Lleras-Muney, 2006 ascribe much of the gains in mortality in the second half of the 20th century to improvements in health technology rather than better direct health inputs such as nutrition or health care. In column 4 of Table 7 we add country specific time trends and find higher calorie consumption still improves mortality outcomes.

In Figure 8 and 9 we plot the country specific fixed effects and time trends from column 4, Tables 6 and 7. Since we adjust for nutrition, we can think of these fixed effects and time trends as representing the disease burden part of health alone. In Figure 8 we plot the fixed effects in infant mortality against the fixed effects in height. As in Figure 6 there is little

evidence of a relationship though people in Sub-Saharan Africa tend to be taller than people elsewhere at each level of disease burden as indicated by infant mortality. One again it appears to be unexplained country fixed factors that explain initial height and infant mortality rather than a common factor, the disease burden.

In Figure 9 we plot the time trends from the regressions in column 4 of Tables 6 and 7. The time trends in height and infant mortality move together, even after correcting for nutrition. While they are correlated the height data suggest declining health due to disease while the infant mortality rate data again suggest improving health. One way of reconciling these results is to argue that the worldwide time trend in infant mortality found in column 3 of Table 7 reflects technological progress in preventing mortality in a given health state. We found no evidence of a time trend in height after adjusting for nutrition in Table 6, which is consistent with no technological progress in turning nutrition into height. If we subtract off the worldwide technological progress estimated in column 3 of Table 7 from the country specific time trends in infant mortality found in the regression in column 4 of Table 7 we get the result plotted in Figure 10. Taking out technological progress in infant mortality, the evidence on the disease burden from trends in infant mortality and trends in height agree quite well across countries.

Our results suggest that we still have omitted variables in levels when we compare heights and infant mortality rates. It may be dangerous to compare across countries using these measures as indicators of health human capital. We leave open the question of what variables would need to be controlled for to make the cross country comparisons feasible. On the other hand, once we control for nutrition and worldwide technological progress in infant mortality, country specific time trends in cohort height and infant mortality rate agree well. Our theory suggests this because, once we have controlled for these additional variables, both measure the same latent variable, the disease burden on children.

We could go further, and add additional control variables, such as income per capita, to our model. However income per capita may be correlated with the provision of medical and public health services that lower the disease burden. Controlling for this will reduce the information content on disease burden in our latent variable. There is a danger of over controlling in equations (5) and (6); we want to control for things that affect height and infant mortality but are not themselves determinants of, or proxies for, the burden of childhood disease.

5. Conclusion

In most of the developing world, and in the historical record of developed countries, there has been a steady picture of advances in infant mortality rates, improvements in nutrition, and increases in adult height, with all of these developments proceeding together. In Sub-Saharan Africa, however, we are seeing a very different pattern unfold. While there have been large reductions in infant mortality, nutrition intake and adult stature have not improved.

The health transition in terms of mortality-morbidity taking place in Sub-Saharan Africa appears to be driven by medical interventions that reduce mortality, rather than by nutrition improvements and broad based reductions in exposure to infectious diseases that would reduce morbidity. This has several implications. It reinforces the view that population health is multidimensional. Movements in mortality measures, such as infant mortality rates and life expectancy, may give a limited picture of how broader population health is changing.

This has implications for studies of the effect of health on worker productivity. In macroeconomic studies of the effect of health on economic growth (Bloom, Canning, and Sevilla 2004) life expectancy or adult mortality rates are often used as measures of population health, assuming that these measures are closely linked to adult height (Shastri and Weil 2003) as used in microeconomic studies (Schultz 2002). This assumption may be unwarranted unless the infant

mortality rate is adjusted to take account of country specific fixed effect and a world wide time trend due to technological progress in mortality reduction.

Our results also call into question the practice of focusing on the use of mortality rates as the sole indicators of population health. For example the Millennium Development Goals (MDGs) focus on reducing mortality, but our results suggest that mortality rates can improve through new technologies that prevent deaths while underlying health problems due to poor nutrition and a high burden of morbidity worsen. Nutrition has been seen as the “forgotten” MDG⁶ and may have benefits birth in terms of reducing infant mortality and in increasing the stature and health human capital on the next generation of adults.

Finally, the continuing child morbidity and lack of physical development may be significant for the future of the aging population in Africa. While the African population is aging (National Research Council 2006), there is strong evidence that the health of adults and the elderly is affected by their childhood health and nutritional status (Fogel and Costa 1997, Catalano and Bruckner 2006, Brush, Harrison, and Waterlow 1997, Blackwell, Hayward, and Crimmins 2001, Yi, Gu, and Land 2007). This suggests that the lack of nutrition and high levels of morbidity among children in Sub-Saharan Africa, and declining adult stature, may be producing unhealthy adults and a growing future health burden.

⁶ World Bank, 2008:

<http://web.worldbank.org/WBSITE/EXTERNAL/NEWS/0,,contentMDK:21627646~pagePK:34370~piPK:34424~theSitePK:4607,00.html>

REFERENCES

- Adetunji, A., and E.R. Bos. 2006. Levels and trends in mortality in Sub-Saharan Africa: Overview. World Bank.
<http://www.ncbi.nlm.nih.gov/bookshelf/br.fcgi?book=dmssa&part=A143>
- Akachi, Y., and D. Canning. 2007. "The Height of Women in Sub-Saharan Africa: the Role of Health, Nutrition, and Income in Childhood." *Annals of Human Biology* 34:397-410.
- Allen, L.H. 1994. Nutritional influences on linear growth: a general review. *Eur J Clin Nutr* 48 Suppl 1:S75-89.
- Alter, G. 2004. "Height, Frailty, and the Standard of Living: Modelling the Effects of Diet and Disease on Declining Mortality and Increasing Height." *Population Studies*, 58: 265–279.
- Barker, D.J.P., P.D. Winter, C. Osmond, B. Margetts, and S.J. Simmonds. 1989. "Weight in Infancy and Death From Ischaemic Heart Disease." *Lancet* 2:577–80.
- Becker, G.S., T.J. Philipson, and R.R. Soares. 2005. "The Quantity and Quality of Life and the Evolution of World Inequality." *American Economic Review* 95: 277–91.
- Blackwell, D., M. Hayward, and E. Crimmins. 2001. "Does Childhood Health Affect Chronic Morbidity Later in Life?" *Social Science and Medicine* 52(8): 1269–84.
- Bloom, D.E., D. Canning, and J.P. Sevilla. 2004. "The Effect of Health on Economic Growth: a Production Function Approach." *World Development* 32: 1–13.
- Bloom D.E., D. Canning. 2000. The health and wealth of nations. *Science* 287(5456): 1207-1209.
- Brush, G., G. Harrison, and J. Waterlow. 1997. "Effects of Early Disease on Later Growth and Early Growth on Later Disease, in Khartoum Infants." *Annals of Human Biology* 24:187–95.
- Catalano, R., and T. Bruckner. 2006. "Child mortality and cohort lifespan: a test of diminished entelechy." *International Journal of Epidemiology* 35:1264–1269.

- Crimmins, E. and C. Finch. 2006. "Infection, Inflammation, Height, and Longevity." *Proceedings of the National Academy of Sciences* 103: 498–503.
- Cutler, D.M., A.S. Deaton, and A. Lleras-Muney. 2006. "The Determinants of Mortality." *Journal of Economic Perspectives* 20: 97–120.
- Deaton, A. 2006. *Global Patterns of Income and Health: Facts, Interpretations, and Policies*. NBER Working Paper No. W12735.
- Deaton, A. 2007. "Height, health, and development." *Proceedings of the National Academy of Sciences* 104: 13232–13237.
- Elo, I.T. and S.H. Preston. 1992. "Effect of Early Life Conditions on Adult Mortality: A Review." *Population Index* 58(2):186–212.
- Fogel, R. 1993. "New Sources and New Techniques for the Study of Secular Trends in Nutritional Status, Health, Mortality and the Process of Aging." *Historical Methods* 26:5–43.
- Fogel, R., and D. Costa. 1997. "A Theory of Technophysio Evolution, with Some Implications for Forecasting Population, Health Care Costs, and Pension Costs." *Demography* 34: 49–66.
- Food and Agricultural Organization. 2006. FAOSTAT data, Food Balance Sheets.
- Greenwood, B.M., A.M. Greenwood, A.K. Bradley, S. Tulloch, R. Hayes, and F.S. Oldfield. 1987. "Deaths in Infancy and Early Childhood in a Well Vaccinated, Rural, West African Population." *Annals of Tropical Paediatrics* 7:91-9.
- Guerrant, R.L., B. Carneiro-Filho, and R.A. Dillingham. 2003. "Cholera, Diarrhea, and Oral Rehydration Therapy: Triumph and Indictment." *Clinical Infectious Diseases* 37:398–405.
- Hayward, M.D. and B.K. Gorman. 2004. "The Long Arm of Childhood: The Influence of Early-Life Social Conditions on Men's Mortality." *Demography* 41:87–107.
- Hoddinott J, Maluccio J, Behrman J, Flores R, Martorell R. 2008. Effect of a nutrition intervention during early childhood on economic productivity in Guatemalan adults. *Lancet* 371(9610): 411-416.

Huffman, S.L., and A. Steel. 1995. "Do Child Survival Interventions Reduce Malnutrition? The Dark Side of Child Survival." In: *Child Growth and Nutrition in Developing Countries*, edited by P. Pinstrip-Anderson, D. Pelletier, and H. Alderman, Ithaca and London: Cornell University Press.

Jacobs, K. and D.A. Sumner. 2002. "The Food Balance Sheets of the Food and Agriculture Organization: A Review of Potential Ways to Broaden the Appropriate Uses of the Data." Food and Agriculture Organization, Rome. 2002. Available at: http://193.43.36.14/abcdq/docs/FBS_Review.pdf

Jacobs, J., T. Katzur, and V. Tassenaar. 2004. On the efficiency of estimators in truncated height samples, University of Groningen, CCSO Centre for Economic Research, Working Paper 200408.

Jamison, D., J. Leslie, and P. Musgrove. 2003. "Malnutrition and Dietary Protein: Evidence from China and from International Comparisons." *Food and Nutrition Bulletin* 24(2):145-54, 56-66.

Komlos, J. 1993. "The Secular Trend in the Biological Standard of Living in the United Kingdom, 1730-1860." *Economic History Review* 46:115-44.

Komlos, J., and B.E. Lauderdale. 2007. "Underperformance in Affluence: The Remarkable Relative Decline in U.S. Heights in the Second Half of the 20th Century." *Social Science Quarterly* 88:283-305.

Kuh, D., R. Hardy, C. Langenberg, M. Richards, and M.E.J. Wadsworth. 2002. "Mortality in Adults Aged 26-54 Years Related to Socioeconomic Conditions in Childhood and Adulthood: Post War Birth Cohort Study." *British Medical Journal* 325:1076-80.

Hegsted, D. 1971. Protein and Calories. FAO/WHO Ad Hoc Committee of Experts on Energy and Protein: Requirements and Recommended Intakes, 22 March-2 April 1971, Rome.

Martorell, R., J.P. Habicht. 1986. Growth in Early Childhood in Developing Countries. In: Falkner, F., J.M. Tanner, editors. *Human Growth Vol. 3*. New York: Plenum.

Martorell, R., A. Lechtig, C. Yarbrough, H. Delgado, R.E. Klein. 1976. Protein-calorie

supplementation and postnatal physical growth: a review of findings from developing countries. *Archivos latinoamericanos de nutrición* 26(2): 115-28.

McKeown, T. 1983. "Food, Infection, and Population." *Journal of Interdisciplinary History* 14: 227-47.

Mendez, M., and L. Adair. 1999. "Severity and Timing of Stunting in the First Two Years of Life Affect Performance on Cognitive Tests in Late Childhood." *Journal of Nutrition* 129:1555-62.

Moradi, A. 2002. "Height and Health of Women in Sub-Saharan Africa and South-Asia 1950-1980." XIII Congress of the International Economic History Association. Buenos Aires, Argentina.

Moradi, A. 2006. "The nutritional status of women in Sub-Saharan Africa, 1950-1980." Department of Economics, University of Oxford.

National Research Council. 2006. "Aging in Sub-Saharan Africa, Recommendations for Furthering Research." Washington, D.C.: National Academies Press.

Peck, M.N., O. Lundberg. 1995. Short stature as an effect of economic and social conditions. *Social Science and Medicine* 41: 733-738.

O'Rand, A.M. and J. Hamil-Luker. 2005. "Processes of Cumulative Adversity: Childhood Disadvantage and Increased Risk of Heart Attack Across the Life Course." *Journals of Gerontology: Psychological Sciences & Social Sciences* 60B (Special Issue II):117-24.

Pinchinat, S., C. Enel , G. Pison , G. Duthé , E. Lagarde , F. Simondon , and K.B. Simondon. 2004. "No Improvement in Weight-for-age of Young Children in Southern Senegal, 1969-1992, Despite a Drastic Reduction in Mortality. Evidence from a Growth Monitoring Programme." *International Journal of Epidemiology* 33:1202-8.

Ruff, C. 2002. "Variation in human body size and shape." *Annual Review of Anthropology* 31:211-32.

Schultz, P. 2002. "Wage Gains Associated with Height as a Form of Health Human Capital." *American Economic Review* 92:349–53.

Schultz, P. 2005. "Productive Benefits of Health: Evidence from Low Income Countries." In: *Health and Economic Growth: Findings and Policy Implications*, edited by G. Lopez-Casasnovas, B. Riveras, and L. Currais. Cambridge, MA: MIT Press.

Shastry, G.K., and D.N. Weil. 2003. "How much of cross-country income variation is explained by health?" *Journal of the European Economic Association* 1:387–96.

Silventoinen K. 2003. Determinants of variation in adult body height. *Journal of Biosocial Science* 2003 Apr; 35(2):263-85. Review.

Silventoinen K, Kaprio J, Lahelma E, Koskenvuo M. 2000. Relative effect of genetic and environmental factors on body height: differences across birth cohorts among Finnish men and women. *American Journal of Public Health* 90, 627-630.

Steckel, R.H. 2009. Heights and human welfare: Recent developments and new directions. *Explorations in Economic History*. 46(1): 1-23.

Steckel, R.H. 2008. "Biological measures of the standard of living." *J Econ Perspect*. 22(1)* 129-152.

Steckel, R. 1995. "Stature and the Standard of Living." *Journal of Economic Literature* 33:1903–40.

Stephensen, C. 1999. "Burden of Infection on Growth Failure." *Journal of Nutrition* 129(2S Supplement):534S–8S.

Stunkard, A.J., T.T. Foch, Z. Hrubec. 1986. A twin study of human obesity. *JAMA* 256(1): 51-4.

Tanner, J.M. 1992. Growth as a measure of the nutritional and hygienic status of a population. *Hormone Research* 38 (supp 1):106-115.

Yi, Z., D. Gu, K.C. Land. 2007. The association of childhood socioeconomic conditions with healthy longevity at the oldest-old ages in China. *Demography* 44(3): 497-516.

Visscher, P., et al. 2006. Assumption-free estimation of heritability from genome-wide identity-by-descent sharing between full siblings. *Public Library of Science Genetics* 24; 2(3): e41.

Weil, D. 2007. "Accounting for the Effect of Health on Economic Growth." *Quarterly Journal of Economics* 122:1265–306.

World Bank. 2005. *World Development Indicators*.

Zerfas, A.J., D.B. Jelliffe, E.F.P. Jelliffe. 1986. Epidemiology and nutrition. In: *Human Growth, a Comprehensive Treatise*, Vol.3. F. Falkner and J.M. Tanner. 2nd Ed. New York: Plenum Press.

Table 1
Descriptive Statistics and Distributional Tests

Birth Cohort	1960	1965	1970	1975
Bangladesh DHS 2004				
Observations	201	285	368	396
Mean height	149.96	150.50	150.33	150.62
Standard Deviation	5.35	5.60	5.60	5.82
Skewness	-0.218	0.372	0.482	0.807
Kurtosis	2.63	4.16	4.24	6.33
Normality test: Shapiro-Wilk p-value	0.17	0.0012	0.0007	<0.00001
Bolivia DHS 2003				
Observations	376	440	496	538
Mean height	150.93	151.29	151.75	151.71
Standard Deviation	5.72	6.38	5.91	5.91
Skewness	-0.03	0.43	0.36	0.26
Kurtosis	3.42	4.29	3.20	4.02
Normality test: Shapiro-Wilk p-value	0.215	0.00004	0.002	0.0005
Ghana DHS 2003				
Observations	102	144	151	187
Mean height	158.59	159.61	160.04	158.57
Standard Deviation	5.36	7.27	5.97	6.92
Skewness	0.23	-2.44	-0.08	-2.49
Kurtosis	2.56	20.90	2.90	21.27
Normality test: Shapiro-Wilk p-value	0.41	0.0001<	0.93	0.0001<

Table 2
Time Trends in Adult Height, Infant Mortality, and Nutrition
Sub-Saharan Africa, 1961- 1985

Country	Adult Height (cm)	Infant Mortality Rate (deaths per 1,000 live birth)	Calories (calories per capita per day)	Protein (grams per capita per day)
Benin	-0.001 (0.014)	-2.310** (0.037)	4.466 (2.486)	0.119 (0.068)
Burkina Faso	0.007 (0.007)	-2.110** (0.023)	0.643 (2.904)	-0.135 (0.088)
Cameroon	0.010 (0.012)	-2.249** (0.014)	8.697** (2.809)	0.008 (0.096)
Central Africa	0.069 (0.042)	-2.905** (0.098)	-2.321 (4.265)	0.180** (0.026)
Chad	-0.034* (0.016)	-3.033** (0.140)	-36.048** (2.845)	-1.360** (0.87)
Cote d'Ivoire	0.024 (0.023)	-3.806** (0.069)	22.622** (2.998)	0.403** (0.074)
Ethiopia	-0.068** (0.017)	-1.784** (0.018)	-3.795 (3.273)	-0.391** (0.109)
Gabon	0.016 (0.032)	-4.365** (0.112)	27.524** (1.725)	1.076** (0.069)
Ghana	0.018 (0.015)	-1.763** (0.026)	-18.089** (5.527)	-0.129 (0.148)
Guinea	-0.021 (0.023)	-2.149** (0.037)	-6.328* (2.119)	-0.051 (0.042)
Kenya	0.050** (0.010)	-2.258** (0.046)	-0.493 (2.248)	-0.319** (0.086)
Madagascar	-0.024 (0.017)	-0.300** (0.0001)	0.118 (1.565)	-0.234** (0.052)
Malawi	-0.005 (0.009)	-2.388** (0.063)	3.552 (3.536)	-0.018 (0.122)

Mali	0.015 (0.015)	-5.431** (0.172)	-13.782** (3.376)	-0.404** (0.090)
Mozambique	-0.011 (0.020)	-1.788** (0.060)	-2.912* (1.284)	-0.155** (0.036)
Niger	-0.015 (0.001)	-0.817** (0.048)	19.282** (2.399)	0.577** (0.107)
Nigeria	-0.010 (0.017)	-0.300** (0.0001)	-12.487** (2.781)	-0.280** (0.083)
Rwanda	-0.023 (0.012)	0.125 (0.071)	18.585** (2.419)	0.230* (0.091)
Senegal	0.074** (0.025)	-2.658** (0.134)	-6.115 (3.287)	0.012 (0.097)
Tanzania	0.005 (0.021)	-1.817** (0.039)	28.325** (3.361)	0.783** (0.081)
Togo	-0.008 (0.025)	-2.578** (0.031)	-9.458** (3.237)	-0.014 (0.081)
Uganda	-0.008 (0.018)	-1.019** (0.093)	-5.479 (2.781)	-0.124 (0.133)
Zambia	0.013 (0.013)	-1.322** (0.092)	-0.272 (2.781)	-0.334** (0.098)
Zimbabwe	-0.009 (0.023)	-1.474** (0.027)	1.673 (2.107)	-0.295** (0.084)

Coefficient of the time trend by country.

Coefficients represent per annum change, standard errors in parentheses, significance level indicated as *(5%), **(1%).

Height trends estimated with weighted least squares; weighted by the number of individuals used to calculate the cohort average height.

Table 3
Time Trends in Adult Height, Infant Mortality, and Nutrition
Non Sub-Saharan African Developing Countries, 1961- 1985

Country	Adult Height (cm)	Infant		
		Mortality Rate (deaths per 1,000 live birth)	Calories (calories per capita per day)	Protein (grams per capita per day)
Bangladesh	0.007 (0.006)	-1.366** (0.094)	-5.519* (2.119)	-0.0007 (0.037)
Bolivia	0.071** (0.005)	-2.262** (0.106)	15.255** (1.981)	0.357** (0.048)
Brazil	0.062** (0.017)	-2.264** (0.019)	18.466** (1.384)	0.217** (0.039)
Colombia	0.096** (0.004)	-2.139** (0.070)	16.721** (2.334)	0.152** (0.004)
Dominican Republic	0.068** (0.001)	-1.656** (0.040)	23.076** (1.055)	0.434** (0.030)
Guatemala	0.040 (0.024)	-2.125** (0.046)	13.117** (2.068)	0.091 (0.046)
Haiti	0.044 (0.020)	-1.939** (0.031)	3.085** (0.905)	0.185** (0.023)
India	0.027** (0.009)	-1.753** (0.038)	6.021* (2.245)	0.054 (0.058)
Indonesia	0.081** (0.010)	-2.396** (0.014)	28.626** (2.028)	0.753** (0.047)
Mexico	0.121** (0.011)	-1.968** (0.033)	37.350** (1.846)	0.950** (0.092)
Morocco	0.085** (0.008)	-1.899** (0.063)	30.844** (1.126)	0.734** (0.065)
Nepal	0.046** (0.010)	-3.747** (0.071)	7.908** (1.557)	0.253** (0.036)
Nicaragua	0.028** (0.005)	-2.464** (0.062)	0.752 (1.619)	-0.331** (0.102)
Peru	0.086**	-2.991**	-2.154	-0.079

	(0.006)	(0.067)	(2.329)	(0.061)
Turkey	0.074**	-3.428**	19.735**	0.383**
	(0.019)	(0.156)	(1.348)	(0.049)

Coefficient of the time trend by country.

Coefficients represent per annum change, standard errors in parentheses, significance level indicated as *(5%), **(1%).

Height trends estimated with weighted least squares; weighted by the number of individuals used to calculate the cohort average height.

Table 4
Regional Time Trends in Adult Height, Infant Mortality, and Nutrition, 1961- 1985

Region	Adult	Infant	Calories	Protein
	Height (cm)	Mortality Rate (deaths per 1,000 live birth)	(calories per capita per day)	(grams per capita per day)
Sub-Saharan Africa	-0.023** (0.003)	-1.711** (0.036)	-0.085 (0.560)	-0.054** (0.017)
Other Developing Countries	0.073** (0.003)	-2.428** (0.027)	13.426** (0.586)	0.294** (0.018)

Coefficient reported on common regional time trend with country fixed effects.

Coefficients represent per annum change, standard errors in parentheses, significance level indicated as *(5%), **(1%).

Height trends estimated with weighted least squares; weighted by the number of individuals used to calculate the cohort average height.

Table 5:
Relationship between Cohort Height and the Infant Mortality Rate in Birth Year
Dependent Variable: cohort average height (centimeters)

	1	2	3	4
Constant	155.339** (0.138)	150.963** (0.100)	149.337** (0.388)	152.012** (0.461)
Log infant mortality rate (per 1,000)	1.516** (0.359)	-1.483** (0.143)	-1.153** (0.171)	-3.557** (0.300)
Energy consumption per capita (calories/100)			0.026 (0.034)	0.017 (0.032)
Protein consumption per capita (grams/100)			2.234* (1.135)	2.191* (1.069)
Time				-0.064** (0.007)
Country fixed effects	no	yes	yes	Yes
R²	0.022	0.977	0.978	0.981
Number of observations	788	788	749	749

Based on data from 39 countries. Weighted least squares where weights are sample size used in calculation of average cohort height. Significance level indicated as *(5%), **(1%).

Table 6:
Relationship Between Cohort Height and Nutrition in Birth Year
Dependent Variable: cohort average height (centimeters)

	1	2	3	4
Constant	157.040** (0.935)	147.789** (0.323)	147.819** (0.325)	151.219** (0.633)
Energy consumption per capita (calories/100)	-0.349** (0.068)	0.133** (0.031)	0.121** (0.034)	-0.042 (0.040)
Protein consumption per capita (grams/100)	10.971 ** (1.830)	-0.029 (1.118)	0.2216 (1.156)	1.4835 (1.226)
Time			0.003 (0.004)	
Country fixed effects	no	no	no	yes
Country fixed effects	no	yes	yes	yes
R²	0.047	0.977	0.977	0.986
N	749	749	749	749

Based on data from 39 countries. Weighted least squares where weights are sample size used in calculation of average cohort height. Significance level indicated as *(5%), **(1%).

Table 7:
Relationship Between Infant Mortality Rate and Nutrition in Birth Year
Dependent Variable: log infant mortality rate

	1	2	3	4
Constant	1.280** (0.061)	1.353** (0.072)	1.268** (0.037)	0.919** (0.037)
Energy consumption per capita (calories/100)	-0.075** (0.004)	-0.083** (0.006)	-0.019** (0.003)	-0.005* (0.002)
Protein consumption per capita (grams/100)	0.788** (0.124)	1.117** (0.209)	-0.220* (0.110)	0.104 (0.076)
Time			-0.018** (0.0002)	
Country fixed effects	no	no	no	Yes
Country fixed effects	no	yes	yes	Yes
R²	0.226	0.709	0.922	0.984
N	1442	1442	1442	1442

Based on data from 39 countries. Ordinary least squares. Significance level indicated as *(5%), ** (1%).

Figure 1




Figure 2




Figure 3

Ghana DHS Comparison




Figure 4

Bolivia DHS Comparison




Figure 5

Egypt DHS Comparison






Figure 6
Estimates of the Level of Child Health
from Infant Mortality Rates and Adult Heights




SSA: Sub-Saharan Africa




SSA: Sub-Saharan Africa

Figure 8
Estimates of the Level of Disease Burden
from Infant Mortality Rates and Adult Heights




SSA: Sub-Saharan Africa



SSA: Sub-Saharan Africa

Figure 10
Estimates of the Time Trend in Disease Burden
Adjusted for Technical Progress in Infant Mortality



SSA: Sub-Saharan Africa